Sewage Treatment System Impact Monitoring Program

Volume 2: Appendices

Data Report 2019-20







Commercial-in-Confidence



Sydney Water

1 Smith Street, Parramatta, NSW Australia 2150 PO Box 399 Parramatta NSW 2124

Report version: STSIMP Data Report 2019-20 Volume 2 final

© Sydney Water 2020

This work is copyright. It may be reproduced for study, research or training purposes subject to the inclusion of an acknowledgement of the source and no commercial usage or sale. Reproduction for purposes other than those listed requires permission from Sydney Water.





Contents



Appendix A: Glossary	1
Appendix B : List of rainfall stations and location details	4
Appendix C : Trends in discharge and reuse volumes, discharge quality and pollufrom coastal WWTPs	
Warriewood WWTP	
North Head WWTP	
Bondi WWTP	
Malabar WWTP	
Cronulla WWTP	
Wollongong WWTP	
Shellharbour WWTP	
Bombo WWTP	65
Malabar storm WWTPs (Fairfield, Glenfield and Liverpool)	72
Appendix D: Trends in discharge and reuse volumes, discharge quality and pollu from inland WWTPs	
Picton WWTP	
West Camden WWTP	
Wallacia WWTP	
Penrith WWTP	_
Winmalee WWTP	
North Richmond WWTP	
Richmond WWTP	
St Marys WWTP	
Quakers Hill WWTP	162
Riverstone WWTP	175
Castle Hill WWTP	184
Rouse Hill WWTP	194
Hornsby Heights WWTP	
West Hornsby WWTP	
Brooklyn WWTP	219
Appendix E : Wastewater overflows	
Dry weather overflows	
Appendix F : Ocean receiving water	
Appendix G : Ocean sediment program	
Appendix H : Harbour and beaches	
Sydney Beaches: Northern Sydney	
Sydney Beaches: Central Sydney	
Sydney Beaches: Southern Sydney	
Illawarra Beaches: Wollongong	
Illawarra Beaches: Shellharbour	
Illawarra Beaches: Bombo	
Sydney Harbour and Estuaries: Botany Bay and Georges River	
Sydney Harbour and Estuaries: Port Hacking	
Sydney Harbour and Estuaries: Port Jackson	
Sydney Harbour and Estuanes. Wildule Harbour	∠00

Sydney Harbour and Estuaries: Pittwater	
Appendix I : Chlorophyll-a trends at estuarine sites	
Port Jackson	
Botany Bay	281
Port Hacking	283
Appendix J : Water quality trends in lagoons	284
Appendix K : Intertidal communities – Shoreline outfalls	292
Intertidal communities Shellharbour 2008 to 2019	296
Appendix L : Intertidal communities of Sydney's estuaries	306
Intertidal rock platform communities	
Settlement panels	309
Appendix M: Hawkesbury-Nepean River water quality trends	311
N92: Nepean River at Maldon Weir	312
N75: Nepean River at Sharpes Weir	318
N67: Nepean River at Wallacia Bridge	324
N57: Nepean River at Penrith Weir	330
N51: Nepean River opposite Fitzgerald Creek	336
N48A: Nepean River at Smith Road	342
N44: Nepean River at Yarramundi Bridge	348
N42: Hawkesbury River at North Richmond	354
N39: Hawkesbury River at Freemans Reach	360
NS04A: Lower South Creek at Fitzroy Bridge	366
N35: Hawkesbury River at Wilberforce	372
NC11A: Lower Cattai Creek at Cattai Ridge Road	378
N3001: Hawkesbury River off Cattai SRA	384
N26: Hawkesbury River at Sackville Ferry	390
N2202: Lower Colo River at Putty Road	
N18: Hawkesbury River at Leets Vale	
NB13: Berowra Creek at Calabash Bay	
NB11: Berowra Creek off Square Bay	414
Appendix N : Stream health of Hawkesbury-Nepean River	
Stream health near Picton WWTP	
Stream health near West Camden WWTP	
Stream health near Wallacia WWTP	
Stream health near Penrith WWTP	
Stream health and community structure of the unnamed creek and the Nepean River near Winmalee WWTP	
Stream health near North Richmond WWTP	461
Stream health near St Mary's WWTP	463
Stream health near Quakers Hill WWTP	464
Stream health near Riverstone WWTP	465
Stream health near Castle Hill WWTP	466
Stream health near Rouse Hill WWTP	467
Stream health near Hornsby Heights WWTP	468
Stream health near West Hornsby WWTP	479
Appendix O: Stream health of other Sydney rivers	492





List of figures

Figure B-1	Rainfall stations used for categorising wastewater data as dry or wet weather days		
Figure B-2	Rainfall stations used for categorising receiving water quality data as dry or wet weather days		
Figure K-1	Barrack Point with a healthy intertidal rock platform community in 2019		
Figure K-2	Barrack Point (in 2001) with an unhealthy intertidal rock platform community impacted by wastewater discharges from the Shellharbour WWTP prior to upgrade in the early to mid 2000's		
Figure K-3	Two-dimensional nMDS ordination plot of 2008 to 2019 intertidal rock platform community data		
Figure K-4	Three-dimensional nMDS ordination plot of 2008 to 2019 intertidal rock platform communidata		
Figure K-5	PCO ordination plot of 2008 to 2019 intertidal rock platform community data -dimensional		
Figure K-6	CAP ordination plot of intertidal rock platform community data (2008 to 2018 for Control site 1 and Control site 2 and 2008 to 2018 outfall site) with 2019 outfall samples (orange squares) predicted		
Figure L-1	Relatively lower salinity zone with year plotted against Principal Coordinates Analysis axis 1 of distance among centroids for sites		
Figure L-2	Relatively higher salinity zone with year plotted against Principal Coordinates Analysis axis 1 of distance among centroids for sites		
Figure L-3	Multiple mean comparison groupings of relatively high salinity locations for 2019-20. (means covered by the same bar are not significantly different)		
Figure L-4	Multiple mean comparison groupings of relatively lower salinity locations for 2019-20. (means covered by the same bar are not significantly different)		
Figure N-1	Stream health of Nepean River near Picton WWTP		
Figure N-2	Stream health of Matahil Creek near West Camden WWTP		
Figure N-3	Stream health of the Nepean River near West Camden WWTP		
Figure N-4	Two-dimensional ordination plot of freshwater macroinvertebrate edge habitat community structure of Matahil Creek upstream and downstream sites of West Camden WWTP		
Figure N-5	Tree diagram of freshwater macroinvertebrate edge habitat community structure of Matah Creek upstream and downstream sites of West Camden WWTP		
Figure N-6	Shade plot of freshwater macroinvertebrate edge habitat community structure of Matahil Creek upstream and downstream sites of West Camden WWTP		
Figure N-7	Dimensions 1 and 2 of three-dimensional ordination plot of freshwater macroinvertebrate edge habitat community structure of sites upstream-downstream of Nepean River at the confluence of Matahil Creek into which West Camden WWTP discharges		
Figure N-8	Tree diagram of freshwater macroinvertebrate edge habitat community structure of sites upstream-downstream of Nepean River at the confluence of Matahil Creek into which We Camden WWTP discharges		
Figure N-9	Shade plot of freshwater macroinvertebrate edge habitat community structure of upstream downstream sites of Nepean River at the confluence of Matahil Creek into which West Camden WWTP discharges		
Figure N-10	Stream health of Warragamba River near Wallacia WWTP		
Figure N-11	Stream health of Boundary Creek near Penrith WWTP		
Figure N-12	Stream health of the Nepean River upstream-downstream of the confluence of Boundary Creek near Penrith WWTP		
Figure N-13	Stream health of unnamed creek below Winmalee WWTP for two downstream sites		
	Stream health of the Nepean River near Winmalee WWTP		
Sewage Tre	eatment System Impact Monitoring Program Vol 2 Data Report 2019-20		

Figure N-15	Dimensions 1 and 3 of three-dimensional ordination plot of freshwater macroinvertebrate edge habitat community structure of both downstream sites of unnamed creek below Winmalee WWTP	
Figure N-16	Dimensions 1 and 2 of three-dimensional ordination plot of freshwater macroinvertebrate riffle habitat community structure of both downstream sites of unnamed creek below Winmalee WWTP	142
Figure N-17	Tree diagram of freshwater macroinvertebrate edge habitat community structure of both downstream sites of unnamed creek below Winmalee WWTP	143
Figure N-18	Tree diagram of freshwater macroinvertebrate riffle habitat community structure of both downstream sites of unnamed creek below Winmalee WWTP	143
Figure N-19	Shade plot of freshwater macroinvertebrate edge habitat community structure of both downstream sites of unnamed creek below Winmalee WWTP	149
Figure N-20	Shade plot of freshwater macroinvertebrate riffle habitat community structure of both downstream sites of unnamed creek below Winmalee WWTP	l50
Figure N-21	Dimensions 1 and 3 of three-dimensional nMDS ordination plot of freshwater macroinvertebrate macrophyte habitat community structure of sites upstream-downstream of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges	152
Figure N-22	Dimensions 1 and 3 of three-dimensional ordination plot of freshwater macroinvertebrate edge habitat community structure of sites upstream-downstream of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges	152
Figure N-23	Tree diagram of freshwater macroinvertebrate macrophyte habitat community structure of sites upstream-downstream of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges	153
Figure N-24	Tree diagram of freshwater macroinvertebrate edge habitat community structure of sites upstream-downstream of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges	153
Figure N-25	Shade plot of freshwater macroinvertebrate macrophyte habitat community structure of upstream-downstream sites of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges	159
Figure N-26	Shade plot of freshwater macroinvertebrate edge habitat community structure of upstream-downstream sites of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges	160
Figure N-27	Stream health of Hawkesbury River upstream-downstream of the confluence of Redbank	161
Figure N-28	Stream health of Redbank Creek near North Richmond WWTP	62
Figure N-29	Stream health of South Creek near St Mary's WWTP	163
Figure N-30	Stream health of Breakfast Creek near Quakers Hill WWTP	164
Figure N-31	Stream health of Eastern Creek near Riverstone WWTP	165
Figure N-32	Stream health of Cattai Creek near Castle Hill WWTP	166
Figure N-33	Stream health of Second Ponds Creek near Rouse Hill WWTP	ŀ67
Figure N-34	Stream health of Calna Creek near Hornsby Heights WWTP	168
Figure N-35	Dimensions 1 and 3 of three-dimensional ordination plot of freshwater macroinvertebrate edge habitat community structure of upstream-downstream sites of Calna Creek near Hornsby Heights WWTP	170
Figure N-36	Dimensions 1 and 2 of three-dimensional ordination plot of freshwater macroinvertebrate riffle habitat community structure of upstream-downstream sites of Calna Creek near Hornsby Heights WWTP	171
Figure N-37	Tree diagram of freshwater macroinvertebrate edge habitat community structure of	ŀ71
Figure N-38	Tree diagram of freshwater macroinvertebrate riffle habitat community structure of upstream-downstream sites of Calna Creek near Hornsby Heights WWTP	172

Figure N-39	Shade plot of freshwater macroinvertebrate edge habitat community structure of upstream-downstream sites of Calna Creek near Hornsby Heights WWTP477	
Figure N-40	Shade plot of freshwater macroinvertebrate riffle habitat community structure of upstream-downstream sites of Calna Creek near Hornsby Heights WWTP	478
Figure N-41	Stream health of Waitara Creek near West Hornsby WWTP	479
Figure N-42	Dimensions 1 and 2 of three-dimensional ordination plot of freshwater macroinvertebrate edge habitat community structure of upstream-downstream sites of Waitara Creek near West Hornsby WWTP	481
Figure N-43	Dimensions 1 and 2 of three-dimensional ordination plot of freshwater macroinvertebrate pool rock habitat community structure of upstream-downstream sites of Waitara Creek near West Hornsby WWTP	481
Figure N-44	Tree diagram of freshwater macroinvertebrate edge habitat community structure of upstream-downstream sites of Waitara Creek near West Hornsby WWTP	482
Figure N-45	Tree diagram of freshwater macroinvertebrate pool rock habitat community structure of upstream-downstream sites of Waitara Creek near West Hornsby WWTP	482
Figure N-46	Shade plot of freshwater macroinvertebrate edge habitat community structure of upstream-downstream sites of Waitara Creek near West Hornsby WWTP	488
Figure N-47	Shade plot of freshwater macroinvertebrate pool rock habitat community structure of upstream-downstream sites of Waitara Creek near West Hornsby WWTP	489
Figure O-1	Stream health of Lane Cove and Parramatta rivers in comparison to control sites	493
Figure O-2	Stream health of lower freshwater Georges River sites compared to control sites in the upper Georges River system	493





List of tables

Table B-1	Rainfall stations used for categorising wastewater data as dry or wet weather days	4
Table B-2	Rainfall stations used for categorising receiving water quality data as dry or wet weather days	6
Table E-1	Trend in dry weather wastewater overflow frequency and volumes for coastal WWTPs wastewater system (2013-14 to 2019-20)	223
Table E-2	Trend in dry weather wastewater overflow frequency and volumes for inland wastewater systems (2013-14 to 2019-20)	224
Table E-3	Trend in wet weather wastewater overflow frequency and volumes for ocean WWTPs wastewater system (2013-14 to 2019-20)	225
Table E-4	Trend in wet weather wastewater overflow frequency and volumes for inland WWTPs wastewater system (2013-14 to 2019-20)	226
Table H-1.	Short-listed dry weather <i>Enterococci</i> exceptions data (>35 cfu/100mL) based on catchment rainfall condition (2019-20)	274
Table K-1	Asymmetrical PERMANOVA of 2019 intertidal assemblages	293
Table K-2	SIMPER 2019 - intertidal assemblages by site	294
Table K-3	Asymmetrical PERMANOVA of 2008 to 2019 intertidal assemblages	300
Table K-4	CAP analysis of 2008 to 2018 intertidal assemblages with 2019 outfall site samples predicted	301
Table N-1	t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from the Nepean River near Picton WWTP	421
Table N-2	t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from Matahil Creek near West Camden WWTP	422
Table N-3	t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from the Nepean River near West Camden WWTP	423
Table N-4	ANOSIM test of 'Site' factor for edge habitat of Matahil Creek upstream and downstream sites of West Camden WWTP	426
Table N-5	PERMANOVA test of 'Site' and 'Year' factors for edge habitat of Matahil Creek upstream and downstream sites of West Camden WWTP	426
Table N-6	PERMDISP test of 'Site' factor for edge habitat of Matahil Creek upstream and downstream sites of West Camden WWTP	426
Table N-7	ANOSIM test of 'Site period' samples for edge habitat of Matahil Creek upstream and downstream sites of West Camden WWTP	427
Table N-8	Genera subset whose multivariate pattern matches full genera set of the edge habitat of Matahil Creek upstream and downstream sites of West Camden WWTP	427
Table N-9	ANOSIM test of Site for edge habitat of upstream-downstream sites of Nepean River at the confluence of Matahil Creek into which West Camden WWTP discharges	431
Table N-10	PERMANOVA test of Site and Year factors for edge habitat of upstream-downstream sites of Nepean River at the confluence of Matahil Creek into which West Camden WWTP discharges	431
Table N-11	PERMDISP test of Site for edge habitat of upstream-downstream sites of Nepean River at the confluence of Matahil Creek into which West Camden WWTP discharges	431
Table N-12	ANOSIM test of Site period for edge habitat of upstream-downstream sites of Nepean River at the confluence of Matahil Creek into which West Camden WWTP discharges	432
Table N-13	Genera subset whose multivariate pattern matches full genera set of the edge habitat of upstream-downstream sites of Nepean River at the confluence of Matahil Creek into which West Camden WWTP discharges	433
Table N-14	t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from the Warragamba River near Wallacia WWTP	435
Table N-15	t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from Boundary Creek near Penrith WWTP	437

Table N-16	t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from the Nepean River near Penrith WWTP437	
Table N-17	t-test of both downstream site SIGNAL-SG scores from 2019-20 for unnamed creek below Winmalee WWTP	439
Table N-18	t-test of upstream-downstream SIGNAL-SG scores from 2019-20 for the Nepean River near Winmalee WWTP	439
Table N-19	ANOSIM test of 'Site' factor for edge habitat unnamed creek near Winmalee WWTP	444
Table N-20	ANOSIM test of 'Site' factor for riffle habitat unnamed creek below Winmalee WWTP	444
Table N-21	PERMANOVA test of 'Site' and 'Year' factors for edge habitat unnamed creek below Winmalee WWTP	445
Table N-22	PERMANOVA test of 'Site' and 'Year' factors for riffle habitat unnamed creek below Winmalee WWTP 445	
Table N-23	PERMDISP test of 'Site' factor for edge habitat unnamed creek below Winmalee WWTP	446
Table N-24	PERMDISP test of 'Site' factor for riffle habitat unnamed creek below Winmalee WWTP	446
Table N-25	ANOSIM test of 'Site period' samples for edge habitat unnamed creek below Winmalee WWTP 447	
Table N-26	ANOSIM test of 'Site period' samples for riffle habitat unnamed creek below Winmalee WWTP	447
Table N-27	Genera subset whose multivariate pattern matches full genera set of the edge habitat unnamed creek below Winmalee WWTP	448
Table N-28	Genera subset whose multivariate pattern matches full genera set of the riffle habitat unnamed creek below Winmalee WWTP	448
Table N-29	ANOSIM test of Site for macrophyte habitat of upstream-downstream sites of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges	454
Table N-30	ANOSIM test of Site for edge habitat Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges	454
Table N-31	PERMANOVA test of Site and Year factors for macrophyte habitat of upstream-downstream sites of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges	455
Table N-32	-	
Table N-33	PERMDISP test of Site for macrophyte habitat of upstream-downstream sites of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges	456
Table N-34	PERMDISP test of Site for edge habitat of upstream-downstream sites of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges	456
Table N-35	ANOSIM test of Site period for macrophyte habitat of upstream-downstream sites of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges	457
Table N-36	ANOSIM test of Site period for edge habitat of upstream-downstream sites of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges	457
Table N-37	Genera subset whose multivariate pattern matches full genera set of the macrophyte habitat of upstream-downstream sites of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges	458
Table N-38	Genera subset whose multivariate pattern matches full genera set of the edge habitat of upstream-downstream sites of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges	458
Table N-39	t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from the Hawkesbury River near North Richmond WWTP	462
Table N-40	t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from Redbank Creek near North Richmond WWTP	462
Table N-41		

Table N-42	t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from Breakfast Creek near Quakers Hill WWTP464	
Table N-43	t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from Eastern Creek near Riverstone WWTP	.465
Table N-44	t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from Cattai Creek near Castle Hill WWTP	.466
Table N-45	t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from Second Ponds Creek near Rouse Hill WWTP	.467
Table N-46	t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from Calna Creek near Hornsby Heights WWTP	.468
Table N-47	ANOSIM test of 'Site' factor for edge habitat of Calna Creek near Hornsby Heights WWTP	.472
Table N-48	ANOSIM test of 'Site' factor for riffle habitat of Calna Creek near Hornsby Heights WWTP	.472
Table N-49	PERMANOVA test of 'Site' and 'Year' factors for edge habitat of Calna Creek near Hornsby Heights WWTP	.473
Table N-50	PERMANOVA test of 'Site' and 'Year' factors for riffle habitat of Calna Creek near Hornsby Heights WWTP	.473
Table N-51	PERMDISP test of 'Site' factor for edge habitat of Calna Creek near Hornsby Heights WWTP	.474
Table N-52	PERMDISP test of 'Site' factor for riffle habitat of Calna Creek near Hornsby Heights WWTP	.474
Table N-53	ANOSIM test of 'Site period' for edge habitat of Calna Creek near Hornsby Heights WWTP	.475
Table N-54	ANOSIM test of 'Site period' for riffle habitat of Calna Creek near Hornsby Heights WWTP	.475
Table N-55	Genera subset whose multivariate pattern matches full genera set of the edge habitat Calna Creek near Hornsby Heights WWTP	.476
Table N-56	Genera subset whose multivariate pattern matches full genera set of the riffle habitat Calna Creek near Hornsby Heights WWTP	.476
Table N-57	t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from Waitara Creek near West Hornsby WWTP	.479
Table N-58	ANOSIM test of 'Site' factor for edge habitat of Waitara Creek near West Hornsby WWTP	.483
Table N-59	ANOSIM test of 'Site' factor for pool rock habitat of Waitara Creek near West Hornsby WWTP	.483
Table N-60	PERMANOVA test of 'Site' and 'Year' factors for edge habitat of Waitara Creek near West	
	Hornsby WWTP	.484
Table N-61	PERMANOVA test of 'Site' and 'Year' factors for pool rock habitat of Waitara Creek near West Hornsby WWTP	.484
Table N-62	PERMDISP test of 'Site' factor for edge habitat of Waitara Creek near West Hornsby WWTP	.485
Table N-63	PERMDISP test of 'Site' factor for pool rock habitat of Waitara Creek near West Hornsby WWTP 485	
Table N-64	ANOSIM test of 'Site period' factor for edge habitat of Waitara Creek near West Hornsby WWTP 486	
Table N-65	ANOSIM test of 'Site period' factor for pool rock habitat of Waitara Creek near West Hornsby WWTP 486	
Table N-66	Genera subset whose multivariate pattern matches full genera set of the edge habitat of Waitara Creek near West Hornsby WWTP	.487
Table N-67	Genera subset whose multivariate pattern matches full genera set of the pool rock habitat of Waitara Creek near West Hornsby WWTP	.487





Appendix A: Glossary

Acronyms/ Abbreviations	Full meanings		
ADCP	Acoustic Doppler Current Profile		
APHA	American Public Health Association		
ANOSIM	Analysis of similarities		
ANOVA	Analysis of variance		
ANZECC	Australian and New Zealand Environment and Conservation Council.		
ANZG	Australian and New Zealand Guidelines for Fresh and Marine Water Quality		
AWI	Antecedent Wetness Index		
AWTP	Advanced Water Treatment Plant		
ВОМ	Bureau of Meteorology		
BOOS	Bondi Ocean Outfall Sewer		
CAP	Canonical Analysis of Principal coordinates		
CBOD	Carbonaceous Biochemical Oxygen Demand		
CCTV	Closed-Circuit Television		
cfu/100mL	Colony forming units per 100 millilitres		
COOS	Cronulla Ocean Outfall Sewer		
CRM	Certified reference material		
CTD	A CTD or Sonde is an oceanography instrument used to measure the conductivity, temperature, and pressure of seawater (the D stands for "depth," which is closely related to pressure		
DF	Degrees of freedom		
DPIE	Department of Planning, Industry and Environment		
DWLP	Dry weather leakage program		
EC ₅₀	Effect Concentration for immobilization of 50% of exposed target biota		
EES	Environment, Energy and Science Branch, NSW DPIE		
EMS	Effluent Management Strategy		
EOP	Emergency Operations Protocol		
EPA	Environment Protection Authority		
EPL	Environment Protection Licence		
ER	Environmental Response		
FOG	Fats, Oils and Grease		
FS&T	Field Sampling and Testing		
GS	Grain Size		
HRC	Healthy River Commission		
IOT	Internet Of Things		
KL	Kilolitre		
km	kilometre(s)		
m	metre		
MOF	Maximum overflow frequency		
Maximum	Maximum value of set of observations		





Acronyms/ Abbreviations	Full meanings		
MCI	Macroinvertebrate Community Index		
MDS	Multidimensional Scaling		
Mean	Mean value of a set of observation		
Median	Median or 50 th percentile value		
mg/L	milligrams per litre		
Minimum	Minimum value of a set of observations		
mL	Millilitre		
ML	Megalitre		
mm	millimetre(s)		
mm³/L	Algal biovolume millimetre cube per litre		
NATA	National Association of Testing Authorities		
NHMRC	National Health and Medical Research Council		
nMDS	Non-metric multidimensional scaling		
No. of Obs	Number of observations		
NSOOS	Northern Suburbs Ocean Outfall Sewer		
NST	Northside Storage Tunnel		
NSW	New South Wales		
NTU	Nephelometric Turbidity unit		
OECD	Organisation for Economic Cooperation and Developments		
ORS	Ocean Reference Station		
OSP	Ocean Sediments Program		
PCA	Principal Component Analysis		
PCB	Polychlorinated Biphenyls		
PCO	Principal Coordinates Ordination		
PCs	Principal Component axes		
PERMANOVA	Permutational Analysis of Variance		
PSR	Pressure-State-Response		
PST	Primary sedimentation tank		
R	Regression co-efficient		
SCAMP	Sewer Catchment Area Management Plan		
SIGNAL-SG	Stream Invertebrate Grade Number Average Level - Genus taxonomic level for the greater Sydney region. Which is a biotic index based on freshwater macroinvertebrate diversity, abundance and tolerance to organic pollution		
SIMPER	Similarity percentage		
SOV	System overflow volume		
SRA	State Recreation Area		
Stats	Statistics		
Std dev	Standard deviation		
STSIMP	Sewage Treatment System Impact Monitoring Program		





Acronyms/ Abbreviations	Full meanings		
SWOOS	Southern and Western Suburbs Ocean Outfall Sewer		
TOC	Total organic carbon		
USEPA	United States Environmental Protection Agency		
WET	Whole Effluent Toxicity		
WWTP	Wastewater treatment plant		
WRP	Water Recycling Plant		
μg/L	micrograms per litre		
μS/cm	micro Siemens per centimetre (unit of conductivity)		





Appendix B : List of rainfall stations and location details

Table B-1 Rainfall stations used for categorising wastewater data as dry or wet weather days

Catchments	Rainfall station (Hydstra code and site name/ description)	Latitude	Longitude	WWTP
Upper Nepean	568053 Picton WWTP	-34.2029	150.6148	Picton and West Camden WWTPs
	568130 West Camden WWTP (composite)	-34.0590	150.6809	
	567163 Regent Ville Rural Fire Service	-33.7745	150.6716	Penrith, St Marys, Glenbrook*, Warragamba* and Wallacia WWTPs
Mid Nepean	567087 St Marys WWTP	-33.7342	150.7692	
	568044 Warragamba Water Filtration Plant	-33.8915	150.5983	- Transgamos and Transgamos
	567084 Quakers Hill WWTP	-33.7365	150.8783	
	567085 Richmond WWTP	-33.6080	150.7671	
Lower Nepean	563069 North Richmond WWTP	-33.5748	150.7156	Quakers Hill, Richmond, North Richmond, Winmalee and Riverstone WWTPs
	563146 Winmalee WWTP	-33.6767	150.6250	
	567100 Riverstone WWTP	-33.6562	150.8477	
	567076 Castle Hill WWTP	-33.7111	150.9842	Castle Hill and Rouse Hill WWTPs
Lower Hawkesbury	567102 Dural (WPS14)	-33.6969	151.0277	
	567120 Brooklyn WWTP	-33.5513	151.1959	
	566055 Hornsby Bowling Club*	-33.7067	151.1070	Brooklyn, West Hornsby and Hornsby
Berowra	566073 Pymble Bowling Club	-33.7408	151.1394	Heights WWTPs
	566053 Hornsby Heights WWTP	-33.6672	151.1047	
	567077 Fairfield WWTP	-33.8807	150.9504	
South West Sydney	567078 Glenfield WWTP	-33.9827	150.9071	Fairfield, Glenfield and Liverpool WWTPs
	566049 Liverpool WWTP	-33.9218	150.9386	







Catchments	Rainfall station (Hydstra code and site name/ description)	Latitude	Longitude	WWTP	
Cronulla	566078 South Cronulla	-34.0700	151.1517	Cremullo WWTD	
	566018 Cronulla WWTP	-34.0307	151.1635	Cronulla WWTP	
Illawarra	568162 Balgownie Reservoir	-34.3928	150.8703		
	568173 Berkeley (Berkeley Sports and Social Club)	-34.4830	150.8473	Bellambi, Port Kembla, Shellharbour, Wollongong and Bombo WWTPs	
	568171 Albion Park Bowling Club	-34.5703	150.7684		
	568181 Figtree Bowling Club	-34.4363	150.8646		
	568188 Kiama Water Tank	-34.6735	150.8434		
North Sydney Coast	566089 Manly Croquet Club (formerly Manly Golf Course)*	-33.7906	151.2758	North Head and Warriewood WWTPs	
	566100 North Head WWTP	-33.8080	151.3019		
	566051 Warriewood WWTP (Composite)	-33.6912	151.2993		
Malabar	566026 Marrickville Bowling Club	-33.9099	151.1641		
	567077 Fairfield WWTP	-33.8807	150.9504	Malabar WWTP	
	567078 Glenfield WWTP	-33.9827	150.9071		
	566049 Liverpool WWTP	-33.9218	150.9386		
Bondi	566032 Paddington (Composite)	-33.8870	151.2253	Bondi WWTP	
	566038 Vaucluse Bowling club	-33.8578	151.2788		

^{*}Not monitored after 2016






Table B-2 Rainfall stations used for categorising receiving water quality data as dry or wet weather days

Catchments	Rainfall station (Hydstra code and site name/ description)	Latitude	Longitude	Water quality monitoring Sites
Upper Nepean	568053 Picton WWTP	-34.2029	150.6148	N92, N75 and N67
	568130 West Camden WWTP (composite)	-34.0590	150.6809	
	568044 Warragamba Water Filtration Plant	-33.8915	150.5983	
Mid Nepean	567163 Regentville Rural Fire Service	-33.7745	150.6716	N57 and N51
	567087 St Marys WWTP	-33.7342	150.7692	
Lower Nepean	567084 Quakers Hill WWTP	-33.7365	150.8783	N48A, N44, N39, N35 and NS04A
	567085 Richmond WWTP	-33.6080	150.7671	
	563069 North Richmond WWTP	-33.5748	150.7156	
	563146 Winmalee WWTP	-33.6767	150.6250	
	567100 Riverstone WWTP	-33.6562	150.8477	
Lower Hawkesbury	567076 Castle Hill WWTP	-33.7111	150.9842	N3001, N26, N18, N2202 and NC11A
	567102 Dural (WPS14)	-33.6969	151.0277	
Berowra	566055 Hornsby Bowling Club*	-33.7067	151.1070	NB13 and NB11
	566073 Pymble Bowling Club	-33.7408	151.1394	
	566053 Hornsby Heights WWTP	-33.6672	151.1047	
Port Jackson Lower	566087 Gladesville Bowling Club	-33.8226	151.1294	PJLC and PJTB
	566073 Pymble Bowling Club	-33.7408	151.1394	
Port Jackson Upper	566087 Gladesville Bowling Club	-33.8226	151.1294	PJ015, PJPRA, PJCB2 and PJDFP
	566082 Auburn RSL Bowling Club	-33.8602	151.0190	
	566032 Paddington (composite)	-33.8870	151.2253	
Middle Harbour	566089 Manly Croquet Club (formerly Manly Golf Course)*	-33.7906	151.2758	PJDR, PJSB and PJCB1
	566100 North Head WWTP	-33.8080	151.3019	





^{*}Not monitored after 2016



Figure B-1 Rainfall stations used for categorising wastewater data as dry or wet weather days







Figure B-2 Rainfall stations used for categorising receiving water quality data as dry or wet weather days

Appendix C: Trends in discharge and reuse volumes, discharge quality and pollutant loads from coastal WWTPs

The coastal WWTPs are presented in the following order from North coast to South coast:

- Warriewood WWTP
- North Head WWTP
- Bondi WWTP
- Malabar WWTP
- Cronulla WWTP
- Wollongong WWTP
- Shellharbour WWTP
- Bombo WWTP
- Malabar storm WWTPs (Fairfield, Glenfield and Liverpool WWTPs) discharging to inland waters in wet weather only

Tests conducted on wastewater are specified under the Environment Protection Licences (EPL) issued by the NSW EPA for each WWTP. All measured analytes that have EPL concentration and load limits are presented in this Appendix.

Under each WWTP, trend plots are presented on:

- Discharge and reuse volume
- Discharge quality
- Discharge loads

Discharge quality and load plots are included in following sub-groups and then analytes presented in alphabetical order:

- Nutrients
- Major conventional analytes
- Toxicity
- Trace metals
- Other chemicals and organics (including pesticides)


Where an analyte shows a trend (positive or negative) two plots are presented (where required) - one with the licence limits and one zoomed in to provide a clearer view of the trend.




Warriewood WWTP

00

Discharge volume and rainfall




Discharge quality: Major conventional analytes











Discharge quality: Trace metals











Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.

Discharge quality: Other chemicals and organics (including pesticides)













Discharge load: Nutrients















North Head WWTP

00

Discharge volume and rainfall




Discharge quality: Major conventional analytes











Discharge quality: Toxicity




Discharge quality: Trace metals











Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.

Discharge quality: Other chemicals and organics (including pesticides)



Note: Statistical test was not conducted because >90% of results were below the detection limit.




Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.










Discharge load: Nutrients











































Discharge load: Other chemicals and organics (including pesticides)







Bondi WWTP

00

Discharge volume and rainfall




Discharge quality: Major conventional analytes











Discharge quality: Toxicity











Discharge quality: Trace metals











Discharge quality: Other chemicals and organics (including pesticides)











Discharge load: Major conventional analytes












Discharge load: Trace metals































Discharge load: Other chemicals and organics (including pesticides)






Note: Discharge loads at Bondi WWTP include discharge loads from network (Diamond Bay and Vaucluse outfalls)




Malabar WWTP

00

Discharge volume and rainfall




Discharge quality: Major conventional analytes











Discharge quality: Toxicity




Discharge quality: Trace metals






Discharge quality: Other chemicals and organics (including pesticides)






Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.











Discharge load: Major conventional analytes












Discharge load: Trace Metals
































Discharge load: Other chemicals and organics (including pesticides)







Cronulla WWTP




Discharge volume and rainfall



Discharge quality: Nutrients




Discharge quality: Major conventional analytes





Note: Statistical test was based on 2019-20 data with previous five years data (2014-2019) due to lab method change in 2014-15.
























Discharge quality: Toxicity













Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.




Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.








detection limit change in 2016-17.




Note: Statistical test was not conducted because >90% of results were below the detection limit.




Note: Statistical test was not conducted because >90% of results were below the detection limit.








Note: Statistical test was not conducted because >90% of results were below the detection limit.




Note: 2019-20 data was compared with previous three years data (2016-2019) due to lab method change in 2016-17. Statistical test was not conducted because >90% of results were below the detection limit.








Discharge load: Major conventional analytes












Discharge load: Trace Metals
































Discharge load: Other chemicals and organics (including pesticides)







Wollongong WWTP




Discharge volume and rainfall











Note: Statistical test was based on 2019-20 data with previous five years data (2014-2019) due to lab method change in 2014-15.




Note: Statistical test was based on 2019-20 data with previous five years data (2014-2019) due to lab method change in 2014-15.











Discharge quality: Trace metals







Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.








Note: Statistical test was not conducted because >90% of results were below the detection limit.




Note: 2019-20 data was compared with previous three years data (2016-2019) due to lab method change in 2016-17. Statistical test was not conducted because >90% of results were below the detection limit.








Discharge load: Major conventional analytes










Note: Discharge loads at Wollongong WWTP also include discharges at Bellambi and Port Kembla WWTPs




Shellharbour WWTP

00


Discharge volume and rainfall




Discharge quality: Nutrients








Note: Statistical test was based on 2019-20 data with previous five years data (2014-2019) due to lab method change in 2014-15.




Note: Statistical test was based on 2019-20 data with previous five years data (2014-2019) due to lab method change in 2014-15.












Discharge quality: Toxicity













Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.

Discharge quality: Other chemicals and organics (including pesticides)











Note: There was a detection limit change in 2016-17. So, statistical test was based on 2019-20 data with previous three years data (2016-2018).












Discharge load: Major conventional analytes















Bombo WWTP




Discharge volume and rainfall











Discharge quality: Major conventional analytes




Note: Statistical test was based on 2019-20 data with previous five years data (2014-2019) due to lab method change in 2014-15.











Discharge quality: Toxicity




Discharge quality: Trace metals











Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.




Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.

Discharge quality: Other chemicals and organics (including pesticides)













Note: Statistical test was based on 2019-20 data with previous three years data (2016-2018) due to detection limit change in 2016-17.












Discharge load: Major conventional analytes












U


Malabar storm WWTPs (Fairfield, Glenfield and Liverpool)



Discharge volume and rainfall (Fairfield)





Discharge quality: Major conventional analytes (Fairfield)




















Discharge quality: Major conventional analytes (Glenfield)




Note: Statistical test was based on 2019-20 data with previous five years data (2014-2019) due to lab method change in 2014-15.











Discharge quality: Major conventional analytes (Liverpool: LP0015 – effluent diversion structure at Chipping Norton)




Note: Statistical test was based on 2019-20 data with previous five years data (2014-2019) due to lab method change in 2014-15.




Note: Statistical test was based on 2019-20 data with previous five years data (2014-2019) due to lab method change in 2014-15.








Discharge quality: Major conventional analytes (Liverpool: LP0076 – recycled water chlorine contact tank)











Discharge quality: Major conventional analytes (Liverpool: LP0081 – overflow chamber downstream of chlorine contact tank)



Note: Statistical test was based on 2019-20 data with previous five years data (2014-2019) due to lab method change in 2014-15.



Appendix D: Trends in discharge and reuse volumes, discharge quality and pollutant loads from inland WWTPs

The inland WWTPs are presented in upstream to downstream locations of the Hawkesbury-Nepean River:

- Picton WWTP
- West Camden WWTP
- Wallacia WWTP
- Penrith WWTP
- Winmalee WWTP
- North Richmond WWTP
- Richmond WWTP
- St Marys WWTP
- Quakers Hill WWTP
- Riverstone WWTP
- Castle Hill WWTP
- Rouse Hill WWTP
- Hornsby Heights WWTP
- West Hornsby WWTP
- Brooklyn WWTP

Tests conducted on wastewater are specified in the Environment Protection Licences (EPL) issued by the NSW EPA for each WWTP. All measured analytes that have EPL concentration and load limits are presented on the following pages.

Under each WWTP, trend plots are presented on:

- Discharge and reuse volume
- Discharge quality
- Discharge loads

Discharge quality and load plots are included in following sub-groups and then analytes presented in alphabetical order:

- Nutrients
- Major conventional analytes
- Toxicity
- Trace metals
- Other chemicals and organics (including pesticides)


Where an analyte shows a trend (positive or negative) two plots are presented (where required) - one with the licence limits and one zoomed in to provide a clearer view of the trend.



Picton WWTP




Discharge volume and rainfall





























Discharge quality: Major conventional analytes (PI0001 Precautionary discharge)




Note: Statistical test was based on 2019-20 data with previous five years data (2014-19) due to lab method change in 2014-15.








Note: Statistical test was based on 2019-20 data with previous five years data (2014-19) due to lab method change in 2014-15.












Discharge quality: Nutrients (PI0011 Irrigation)




















Discharge quality: Major conventional analytes (PI0011 Irrigation)





Note: Note: Statistical test was based on 2019-20 data with previous five years data (2014-19) due to lab method change in 2014-15

































Discharge quality: Major conventional analytes (PI0013 Irrigation)
















Discharge load: Nutrients












Discharge load: Major conventional analytes















West Camden WWTP



Discharge volume and rainfall










Discharge quality: Nutrients

















Discharge quality: Major conventional analytes

(Source=Model: DF=1:

F Value=4.4713,

Pr>F=0.0349)





Financial year


Note: 2019-20 data was compared with previous five years data (2014-2019) due to lab method change in 2014-15. Statistical test was not conducted because >90% of results were below the detection limit.















Discharge quality: Toxicity







Discharge quality: Trace metals













Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.




Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.








Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.



Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.




Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.











Note: Statistical test was not conducted because >90% of results were below the detection limit.




Note: Statistical test was not conducted because >90% of results were below the detection limit.




Note: 2019-20 data was compared with previous three years data (2016-2019) due to lab method change in 2016-17. Statistical test was not conducted because >90% of results were below the detection limit.



Discharge load: Nutrients





Discharge load: Major conventional analytes















Wallacia WWTP



Discharge volume and rainfall





Discharge quality: Nutrients

























Note: Statistical test was based on 2019-20 data with previous five years data (2014-2019) due to lab method change in 2014-15.









Discharge quality: Toxicity






Note: Statistical test was not conducted because >90% of results were recorded at 100% survival for Ceriodaphnia dubia

Discharge quality: Trace metals











Discharge quality: Other chemicals and organics (including pesticides)



Note: 2019-20 data was compared with previous three years data (2016-2019) due to lab method change in 2016-17.






Discharge load: Nutrients

900

600

300

2010-11



Discharge load: Major conventional analytes


2011-12

2012-13

2013-14

2014-15

Financial year



2015-16

2016-17

2017-18

2018-19

240


120


60

2019-20












Penrith WWTP



Discharge volume and rainfall







2010-11


2011-12

(Source=Model; DF=1; F Value=0.4578;

2012-13


2013-14

Pr>F=0.4989)




2014-15

Financial year



2015-16

2016-17




2018-19

2019-20


2017-18








Discharge quality: Major conventional analytes




Note: Statistical test was based on 2019-20 data with previous five years data (2014-2019) due to lab method change in 2014-15.




Note: Monitoring of total chlorine residual started in January 2015.










Discharge quality: Toxicity





Note: Statistical test was not conducted because >90% of results were recorded at 100% survival for Ceriodaphnia dubia.

















Note: 2019-20 data was compared with previous three years data (2016-2019) due to lab method change in 2016-17. Statistical test was not conducted because >90% of results were below the detection limit.




















Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.




Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.











Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.



Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.

Discharge quality: Other chemicals and organics (including pesticides)




Note: 2019-20 data was compared with previous three years data (2016-2019) due to lab method change in 2016-17. Statistical test was not conducted because >90% of results were below the detection limit.




Discharge load: Nutrients






Discharge load: Major conventional analytes












Discharge load: Trace metals






























Discharge load: Other chemical and organics (including pesticides)







Winmalee WWTP



Discharge volume and rainfall





Discharge quality: Nutrients























Discharge quality: Major conventional analytes




Note: Statistical test was based on 2019-20 data with previous five years data (2014-2019) due to lab method change in 2014-15.










Discharge quality: Toxicity


























Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.




Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.











Discharge quality: Other chemicals and organics (including pesticides)




Note: Statistical test was not conducted because >90% of results were below the detection limit.



Discharge load: Nutrients





Discharge load: Major conventional analytes















North Richmond WWTP



Discharge volume and rainfall





Discharge quality: Nutrients























Discharge quality: Major conventional analytes




Note: Statistical test was based on 2019-20 data with previous five years data (2014-2019) due to lab method change in 2014-15.











Discharge quality: Toxicity


























Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.




Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.







Discharge quality: Other chemicals and organics (including pesticides)



Note: Statistical test was not conducted because >90% of results were below the detection limit.




Note: Statistical test was not conducted because >90% of results were below the detection limit.




Note: 2019-20 data was compared with previous three years data (2016-2019) due to lab method change in 2016-17. Statistical test was not conducted because >90% of results were below the detection limit.









Discharge load: Major conventional analytes















Richmond WWTP



Discharge volume and rainfall





























Discharge quality: Major conventional analytes (RM0016 Bypass Effluent)





Note: 2019-20 data was compared with previous five years data (2014-2019) due to lab method change in 2014-15. Statistical test was not conducted because >90% of results were below the detection limit.


















Discharge quality: Toxicity (RM0016 Bypass Effluent)






Note: Statistical test was not conducted because >90% of results were recorded at 100% survival for Ceriodaphnia dubia.

Discharge quality: Nutrients (RM0017 Effluent)











Discharge quality: Major conventional analytes (RM0017 Effluent)





2014-15.

























Discharge load: Nutrients




Discharge load: Major conventional analytes




Financial year













St Marys WWTP



Discharge volume and rainfall






























Discharge quality: Major conventional analytes



















Discharge quality: Toxicity






Discharge quality: Trace metals























Discharge quality: Other chemicals and organics (including pesticides)



Note: Statistical test was not conducted because >90% of results were below the detection limit.




Note: 2019-20 data was compared with previous three years data (2016-2019) due to lab method change in 2016-17. Statistical test was not conducted because >90% of results were below the detection limit.




Discharge load: Nutrients






Discharge load: Major conventional analytes












Discharge load: Trace metals






































Quakers Hill WWTP



Discharge volume and rainfall










Discharge quality: Nutrients







Discharge quality: Major conventional analytes






Note: 2019-20 data was compared with previous five years data (2014-2019) due to lab method change in 2014-15. Statistical test was not conducted because >90% of results were below the detection limit.











Discharge quality: Toxicity





Note: Statistical test was not conducted because >90% of results were recorded at 100% survival for Ceriodaphnia dubia.














Note: Statistical test was not conducted because >90% of results were below the detection limit.


























Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.














Discharge quality: Other chemicals and organics (including pesticides)




Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17. Statistical test was not conducted because >90% of results were below the detection limit.


Discharge load: Nutrients
















































Discharge load: Other chemicals and organics (including pesticides)











Discharge volume and rainfall





Discharge quality: Nutrients























Discharge quality: Major conventional analytes




Note: Statistical test was based on 2019-20 data with previous five years data (2014-2019) due to lab method change in 2014-15.











Discharge quality: Toxicity


























Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.




Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.




Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.








Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.




Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.

Discharge quality: Other chemicals and organics (including pesticides)











Note: 2019-20 data was compared with previous three years data (2016-2019) due to lab method change in 2016-17. Statistical test was not conducted because >90% of results were below the detection limit.


Discharge load: Nutrients


















Castle Hill WWTP



Discharge volume and rainfall












Discharge quality: Nutrients















Discharge quality: Major conventional analytes




Note: Statistical test was based on 2019-20 data with previous five years data (2014-2019) due to lab method change in 2014-15.











Discharge quality: Toxicity




Note: Statistical test was not conducted because >90% of results were recorded at 100% survival for *Ceriodaphnia dubia*.

Discharge quality: Trace metals












Note: Statistical test was not conducted because >90% of results were below the detection limit.













Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.




Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.




Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.








Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.




Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.

Discharge quality: Other chemicals and organics (including pesticides)










Note: 2019-20 data was compared with previous three years data (2016-2019) due to lab method change in 2016-17. Statistical test was not conducted because >90% of results were below the detection limit.



Discharge load: Nutrients



Discharge load: Major conventional analytes

2011-12

2012-13

2013-14

201415

Financial year

2015-16


2016-17

2017-18

2018-19


2019-20


2010-11















Rouse Hill WWTP



Discharge volume and rainfall






























Discharge quality: Major conventional analytes (RH0004 Effluent)

















Discharge quality: Toxicity (RH0004 Effluent)







Discharge quality: Trace metals (RH0004 Effluent)













Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.




Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.








Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.




Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.

Discharge quality: Other chemicals and organics (including pesticides) (RH0004 Effluent)






Discharge load: Nutrients




Discharge load: Major conventional analytes




Financial year













Hornsby Heights WWTP



Discharge volume and rainfall





Discharge quality: Nutrients

























Note: Statistical test was based on 2019-20 data with previous five years data (2014-2019) due to lab method change in 2014-15.









Discharge quality: Toxicity






Note: Statistical test was not conducted because >90% of results were recorded at 100% survival for Ceriodaphnia dubia.


Discharge quality: Trace metals













Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.




Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.








Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.




Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.

Discharge quality: Other chemicals and organics (including pesticides)











Note: Statistical test was not conducted because >90% of results were below the detection limit.



Note: 2019-20 data was compared with previous three years data (2016-2019) due to lab method change in 2016-17. Statistical test was not conducted because >90% of results were below the detection limit.






2015-16

2016-17

2017-18

2018-19

2019-20




Discharge load: Major conventional analytes

2010-11


2011-12


2012-13















West Hornsby WWTP

00

Discharge volume and rainfall




Discharge quality: Nutrients











Discharge quality: Major conventional analytes




Note: Statistical test was based on 2019-20 data with previous five years data (2014-2019) due to lab method change in 2014-15.











Discharge quality: Toxicity




Note: Statistical test was not conducted because >90% of results were recorded at 100% survival for *Ceriodaphnia dubia*.

Discharge quality: Trace metals











Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.



Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.




Note: Statistical test was based on 2019-20 data with previous three years data (2016-2019) due to detection limit change in 2016-17.






Discharge quality: Other chemicals and organics (including pesticides)




Note: Statistical test was not conducted because >90% of results were below the detection limit.




Note: 2019-20 data was compared with previous three years data (2016-2019) due to lab method change in 2016-17. Statistical test was not conducted because >90% of results were below the detection limit.



Discharge load: Nutrients





Discharge load Major conventional analytes















Brooklyn WWTP



Discharge volume and rainfall




Discharge quality: Nutrients












Discharge quality: Major conventional analytes























Note: Statistical test was not conducted because >90% of results were recorded at 100% survival for Ceriodaphnia dubia.







Dry weather overflows

Table E-1 Trend in dry weather wastewater overflow frequency and volumes for coastal WWTPs wastewater system (2013-14 to 2019-20)

Wastewater	2013-	-14	2014	-15	2015-	16	2016-	17	2017-	18	2018-	19	2019-20	
system	Frequency	Volume (KL)												
Warriewood	52	309	34	272	18	62	6	99	6	39	3	27	7	55
North Head / Northern suburbs	441	8,331	406	9,248	241	7,379	98	6,547	147	10,197	170	16,151	176	7,948
Bondi	18	2,154	12	1,208	12	448	24	1,017	16	960	30	1,424	22	1,480
Malabar/Southern suburbs	91	7,837	100	13,997	77	6,803	76	8,098	75	6,112	79	6,853	133	9,530
Cronulla	25	289	26	795	19	424	30	1,030	42	2,205	54	2,279	41	693
Wollongong	17	126	11	188	16	472	9	174	11	132	28	551	27	649
Port Kembla	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Shellharbour	3	8	1	50	4	75	2	75	3	387	3	42	4	172
Kiama/Bombo	2	25	2	13	1	2	1	142	1	7	2	34	2	39
All ocean systems	649	19,080	592	25,771	388	15,664	246	17,182	301	20,039	369	27,361	412	20,567



Table E-2 Trend in dry weather wastewater overflow frequency and volumes for inland wastewater systems (2013-14 to 2019-20)

Wastewater	2013	-14	2014	-15	2015-	16	2016-	17	2017-	18	2018	2018-19 2019		
system	Frequency	Volume (KL)												
Picton	4	171	0	0	1	83	5	188	2	22	3	45	4	28
West Camden	5	18	2	16	2	43	4	35	7	1,079	1	7	3	35
Wallacia	0	0	0	0	1	24	2	3	1	13	1	9	0	0
Penrith	13	896	15	483	9	492	6	194	11	287	3	73	10	180
Winmalee	11	250	13	263	13	142	11	441	11	580	8	180	5	99
North Richmond	0	0	0	0	0	0	0	0	0	0	2	14	1	6
Richmond	0	0	0	0	1	0.3	0	0	0	0	0	0	0	0
St Marys	5	1,823	4	151	3	235	2	62	0	0	4	170	8	192
Quakers Hill	38	298	28	153	21	90	2	109	5	123	10	866	4	130
Riverstone	1	11	0	0	1	0.7	0	0	1	87	1	57	2	36
Castle Hill	12	394	9	253	10	342	2	312	4	74	8	213	2	20
Rouse Hill	14	162	10	55	6	32	6	72	1	10	9	318	8	163
Hornsby Heights	19	156	15	81	16	95	6	43	3	2	4	37	9	147
West Hornsby	37	1,186	32	688	13	185	8	123	6	27	9	391	5	100
Brooklyn-Danger Island	0	0	0	0	0	0	0	0	0	0	0	0	0	0

1,764

97

54

1,582

52

2,304

63

2,380

61



5,365

2,144

128

All inland

systems

159

1,138



2013-14 to 2010-

Table E-3 Trend in wet weather wastewater overflow frequency and volumes for ocean WWTPs wastewater system (2013-14 to 2019-20)

Wastawatar ayatam	20)13-14	20)14-15	20)15-16	20)16-17	20	017-18	2018-19		2019-20	
Wastewater system	MOF	SOV (ML)	MOF	SOV (ML)	MOF	SOV (ML)								
Warriewood	1	0	2	3	3	159	3	0.5	1	3.5	2	4.8	5	157.7
North Head / Northern suburbs	11	666.7	23	11,181	26	9,076	23	4,000	16	279.9	23	3,801.0	16	9,861.0
Bondi	15	10.6	23	189	23	247	20	94	5	11.7	10	179.5	12	489.1
Malabar/Southern suburbs	14	1303.5	33	15,098	28	10,515	29	5,328	20	2,415.0	28	6,586.5	18	15,593.2
Cronulla	1	0	13	815	9	678	13	94	-	0.03	8	28.0	9	659.7
Wollongong	6	126.5	21	285	6	367	8	86	2	0.2	5	25.0	2	59.3
Bellambi	14	315.7	6	45	14	693	22	234	1	0.0	19	46.2	4	159.8
Port Kembla	6	143.1	7	78	4	503	8	159	2	0.7	4	6.7	2	142.4
Shellharbour	4	121.4	6	223	4	494	4	146	1	1.5	4	2.6	1	106.3
Kiama/Bombo	6	29.1	9	47	4	216	8	72	4	2.5	6	4.1	2	3.0
All ocean systems	78	2,717	143	27,964	121	22,949	138	10,213.5	52	2,715.0	109	10,684.4	71	27,232

MOF: Maximum overflow frequency

SOV: System overflow volume

U

00

Table E-4 Trend in wet weather wastewater overflow frequency and volumes for inland WWTPs wastewater system (2013-14 to 2019-20)

Mastaurator avatam	20 ⁻	2013-14		14-15	2015-16		2016-17		2017-18		2018-19		2019-20	
Wastewater system	MOF	SOV (ML)	MOF	SOV (ML)	MOF	SOV (ML)	MOF	SOV (ML)	MOF	SOV (ML)	MOF	SOV (ML)	MOF	SOV (ML)
Picton	0	0	0	0	1	8	0	0	0	0	0	0	1	7.5
West Camden	1	0.4	3	32	3	211	3	3	0	0	2	1.0	1	65.1
Wallacia	0	0	3	14	5	11	5	11	0	0	3	2.2	4	28.6
Penrith	1	0	3	27	5	22	4	0.3	0	0	4	12.2	6	173.0
Winmalee	0	0	2	0.1	0	0	0	0	0	0	1	0.1	2	98.0
North Richmond	0	0	2	13	2	17	8	11	0	0	2	0.4	3	15.6
Richmond	0	0	2	1.2	0	0	0	0	0	0	0	0	1	2.1
St Marys	3	8.2	10	222	5	400	5	38	0	0	10	71.7	6	399.7
Quakers Hill	4	20.4	10	744	7	693	10	162	1	12.2	8	280.0	4	538.2
Riverstone	1	0.0	1	12	2	30	3	3.5	0	0	2	0.5	1	34.9
Castle Hill	1	0.4	2	49	2	32	2	1	0	0	4	4.6	2	75.5
Rouse Hill	0	0	1	140	3	94	3	0.3	0	0	2	8.1	1	111.9
Hornsby Heights	1	0.03	1	0.4	1	1	0	0	0	0	0	0	1	1.1
West Hornsby	1	0.03	5	125	5	162	2	11	0	0	8	42.9	2	91.8
Brooklyn-Danger Island	0	0.0	0	0	0	0	0	0	0	0	0	0	0	0.0
All inland systems	13	29.5	45	1,380	41	1,680	45	241	1	12.2	46	423.7	15	1,643





Appendix F: Ocean receiving water

The 2019-2020 summary of modelled chemical concentrations near deepwater ocean outfalls is presented in the separate report titled the 2020 Ocean Sediment Program Assessment Report. Context for that separate report is outlined in Appendix G.







Appendix G : Ocean sediment program

See the separate report titled the 2020 Ocean Sediment Program Assessment Report. Monitoring under the Ocean Sediment Program enables assessment of the longer-term performance of the deepwater ocean outfalls and provide a mechanism to alert for development of possible long-term accumulative effects. The 2020 report analyses previously collected assessment data from the years of 2002, 2005, 2008, 2011, 2014, and 2016 along with relatively recently collected 2020 data.





Appendix H: Harbour and beaches

The analysis of the Beachwatch data has been designed to identify potential wastewater overflows or leakage under dry weather conditions. Overflows or leakage reaching the waterways during dry weather conditions pose a risk to public health. The wet weather public health risk for recreational activities in waterways (harbour and beaches) are well known.

Assumptions behind the data for Beachwatch analysis:

- Enterococci results without a respective conductivity value were excluded. Conductivity results for many sites were not available prior to 2013. Conductivity data is required to separate dry weather data from wet weather data.
- Only dry weather results were included in these plots. *Enterococci* results associated with conductivity below 30,000 μ S/cm were considered wet weather and not included in these plots
- Data labels are shown in plots for all *Enterococci* values ≥ 230 cfu/100mL, which is the secondary contact recreation guideline (ANZECC 2000).

The Beachwatch results are presented in the following order similar to monitoring programs and sub-catchments as stated in the method section of Volume 1:

Sydney Beaches

- Northern Sydney
- Central Sydney
- Southern Sydney

Illawarra Beaches

- Wollongong
- Shellharbour
- Bombo

Harbours

- Botany Bay and Georges River
- Port Hackings
- Port Jackson
- Middle Harbour
- Pittwater

The sites under each sub-catchment are presented in the order from north coast to south coast. When the sub-catchment is a harbour with sites on both coasts then sites on south coasts were stated first and then following clockwise direction to the north coast.


Note: The size of the bubble can't be compared between plots precisely due to software limitations. The bubble size for each site varied based on maximum value for that site using the following ranges, which helped in adjusting the size of bubble for comparison between plots.


Bubble size	Range of maximum Enterococci value (cfu/100mL)
5	<200
6	200 to 399
7	400 to 599
8	600 to 799
9	800 to 999
10	1,000 to 2,999
11	3,000 to 5,999
12	6,000 to 9999
15	10,000 or more








Sydney Beaches: Northern Sydney

















35,000

30,000 4 30JUN10

30JUN11

30JUN12

30JUN13

30JUN14

30JUN15


30JUN16


30JUN17

30JUN18

30JUN19


30JUN20


















Sydney Beaches: Central Sydney


















































Sydney Beaches: Southern Sydney

























Illawarra Beaches: Wollongong

30JUN10

30JUN11

30JUN12

30JUN13

30JUN14

30JUN15


Sampling date

30JUN16

30JUN17


30JUN18


30JUN19













































00

Illawarra Beaches: Shellharbour











Illawarra Beaches: Bombo
















Sydney Harbour and Estuaries: Botany Bay and Georges River



































































Sydney Harbour and Estuaries: Port Hacking





















Sydney Harbour and Estuaries: Port Jackson


































Sampling date

















Sydney Harbour and Estuaries: Middle Harbour
















30JUN13

30JUN10

30JUN11

30JUN12

30JUN14

30JUN15

Sampling date

30JUN16


30JUN17

30JUN18

30JUN19







30JUN13

30JUN14

30JUN15

Sampling date

30JUN16

30JUN17

30JUN18

30JUN19

40,000

35,000

30JUN10


30JUN11

30JUN12






Sydney Harbour and Estuaries: Pittwater









30JUN13

30JUN14

30JUN15

Sampling date

30JUN16

30JUN17

30JUN18

30JUN19


30JUN11


30JUN10


30JUN12













40,000

35,000


30,000

30JUN10

30JUN11

30JUN12





30JUN14

30JUN13

30JUN15

Sampling date

30JUN16

30JUN17

30JUN18

30JUN19





Table H-1. Short-listed dry weather *Enterococci* exceptions data (>35 cfu/100mL) based on catchment rainfall condition (2019-20)

Sampling site	Sampling date	Enterococci (cfu/100ML)	Conductivity (μS/cm)	72 hours rain ≤2 mm	Station no	Station name
Avalon Beach	18-Oct-19	130	53900	0	66079	Avalon Beach (Palmgrove Rd))
Balmoral Baths	24-Jan-20	130	54200	1.4	66006	Sydney Botanic Garden
Barrenjoey Beach	8-Oct-19	270	52200	0	66128	Palm Beach (Sunrise Road)
	20-Mar-20	260	53500	1.2	68228	Bellambi AWS
Bellambi Beach	1-Apr-20	2500	31800	0.8		
	9-Aug-19	100	51100	0	66126	Collaroy (Long Reef Golf Club)
Bilarong Reserve	16-Jan-20	1500	45500	0		
	29-Jan-20	62	42700	0		
	22-Apr-20	53	47800	0		
	25-May-20	150	50700	1.4		
	4-Sep-19	130	52200	0		
Boat Harbour	10-Oct-19	140	53900	2	00050	One One (Public Only)
	13-Jan-20	35	54500	0	66058	Sans Souci (Public School
	23-Mar-20	48	54000	1		
Bombo Beach	15-Dec-19	39	54200	0.2	68242	Kiama (Bombo Headland)
Bondi Beach	22-Oct-19	810	53900	0.9	66098	Rose Bay (Royal Sydney Golf Club)
Brighton Le Sands Baths	24-Feb-20	150	43800	1.6	66037	Sydney Airport AMO
Duranta Danah	12-Dec-19	46	54700	0	66098	Rose Bay (Royal Sydney Golf Club)
Bronte Beach	24-Jan-20	38	54200	0		
Bulli Beach	1-Apr-20	570	50500	0.8	68108	Bellambi AWS
Camp Cove	12-Dec-19	89	54500	0	66098	Rose Bay (Royal Sydney Golf Club)
Obio and Danah	19-Nov-19	110	53400	0	66006	Sydney Botanic Garden
Chinamans Beach	24-Jan-20	58	53900	1.4		
Clareville Beach	7-May-20	44	54200	0.6	66079	Avalon Beach (Palmgrove Rd))
Clifton Gardens	24-Jan-20	64	54200	1.4	66006	Sydney Botanic Garden
Clontarf Pool	27-Apr-20	94	53700	0	66080	Castle Cove (Rosebridge Ave)
Clavelly Basel	4-Sep-19	300	53700	0	66052	Randwick (Randwick St)
Clovelly Beach	23-Apr-20	37	55500	0	66052	Randwick (Randwick St)
Collaroy Beach	9-Aug-19	130	54300	0	66126	Collaroy (Long Reef Golf Club)
Como Baths	10-Oct-19	83	45600	2	66058	Sans Souci (Public School
	22-Oct-19	74	53700	0	66037	Sydney Airport AMO
Congwong Bay	12-Dec-19	61	53800	0		
	24-Jan-20	140	54100	0		
Coogee Beach	4-Sep-19	70	53900	0	66098	Rose Bay (Royal Sydney Golf Club)
	12-Dec-19	170	54600	0		
	24-Jan-20	140	54000	0		
	4-May-20	66	54100	0		
Corrimal Beach	18-Jun-20	55	52900	0.4	68228	Bellambi AWS
Davidson Reserve	20-Mar-20	44	40300	0	66188	Belrose (Evelyn Place)









Sampling site	Sampling date	Enterococci (cfu/100ML)	Conductivity (μS/cm)	72 hours rain ≤2 mm	Station no	Station name
Little Bay Beach	13-Nov-19	38	54700	0	66052	Randwick (Randwick St)
	12-Dec-19	50	53600	0		
	24-Jan-20	390	54200	0.1		
Little Manly Cove	12-Dec-19	200	54400	0	66126	Collaroy (Long Reef Golf Club)
Long Reef Beach	6-Sep-19	83	54000	0		Collaroy (Long Reef Golf Club)
	13-Aug-19	150	54800	0.6		Randwick (Randwick St)
	13-Nov-19	55	54500	0		
	12-Dec-19	94	54200	0		
Malabar Beach	24-Jan-20	250	54200	0.1	66052	
	23-Apr-20	38	55500	0		
	4-May-20	81	54800	0		
	25-Jun-20	35	54700	1		
	12-Dec-19	82	54600	0		
Manly Cove	18-Dec-19	40	54200	0	66126	O-11 (1 D
Marily Cove	30-Dec-19	63	54400	0	66126	Collaroy (Long Reef Golf Club)
	4-May-20	80	54300	0		
	24-Jan-20	330	54600	0.1	66053	
Maroubra Beach	23-Apr-20	260	54900	0		Randwick (Randwick St)
	4-May-20	58	54600	0		
Mana Vala Basah	9-Aug-19	120	54100	0	66079	Avalon Beach (Palmgrove Rd))
Mona Vale Beach	18-Oct-19	35	54800	0.4	66141	Mona Vale Golf Club
Monterey Baths	28-Jan-20	66	53100	0	66058	Sans Souci (Public School
Murray Rose Pool	4-May-20	40	53000	0	66098	Rose Bay (Royal Sydney Golf Club)
	9-Aug-19	80	51000	0.4	66141	Mona Vale Golf Club
	18-Nov-19	79	43400	0	66141	
Narrabeen Lagoon (Birdwood	30-Dec-19	35	45800	0		
Park)	22-Apr-20	120	51900	0.2		
	5-Jun-20	35	52100	0.2		
	18-Jun-20	36	49800	0.2		
Newport Beach	9-Aug-19	39	53700	0	66079	Avalon Beach (Palmgrove Rd))
Nielsen Park	24-Jan-20	37	54000	0	66098	Rose Bay (Royal Sydney Golf Club)
North Curl Curl Beach	9-Aug-19	60	54800	0	66126	Collaroy (Long Reef Golf Club)
	9-Aug-19	180	54200	0.4	66141 Mona Vale Golf Club	
North Narrabeen Beach	6-Sep-19	36	50100	0.6		Mona Vale Golf Club
	16-Jan-20	100	54100	0.2		
North Wollongong Beach	14-Jan-20	780	54200	0	68228	Bellambi AWS
Northbridge Baths	24-Jan-20	46	52800	0	66080	Castle Cove (Rosebridge Ave)
Paradise Beach Baths	18-Nov-19	260	54000	0	66128	Palm Beach (Sunrise Road)
	27-Nov-19	46	54400	0.8	68131	Port Kembla (BSL Central Lab)






Sampling site	Sampling date	Enterococci (cfu/100ML)	Conductivity (μS/cm)	72 hours rain ≤2 mm	Station no	Station name
Port Kembla Beach	9-Dec-19	39	54100	0		
	15-Dec-19	70	54200	0		
	14-Mar-20	210	53700	1.2	68110	Berkeley (Northcliffe Drive)
	25-May-20	47	54600	1.2		
Ramsgate Baths	28-Jan-20	130	52900	0	66058	Sans Souci (Public School
Rose Bay Beach	4-May-20	140	53800	0	66098	Rose Bay (Royal Sydney Golf Club)
Shelly Beach (Manly)	9-Aug-19	94	55000	0	66126	Collaroy (Long Reef Golf Club)
	16-Jan-20	100	54400	0		
Cilven D I	10-Oct-19	60	53400	2	66058	Sans Souci (Public School
Silver Beach	11-Nov-19	100	53600	0	00038	Sans Souci (Fubile School
	6-Sep-19	53	54100	0		Oallana (Laran Baat Oalt Olah)
South Curl Curl Beach	15-Oct-19	39	54200	1	66126	
	9-Dec-19	60	54400	0	00120	Collaroy (Long Reef Golf Club)
	13-Jan-20	53	54500	1.4		
	23-Apr-20	230	55000	0		
	4-May-20	59	54300	0		Randwick (Randwick St) Collaroy (Long Reef Golf Club)
South Maroubra Beach	15-Jul-19	50	54400	0	66052	
Beach	23-Apr-20	120	54900	0		
	4-May-20	45	55000	0		
	12-Jul-19	43	54400	0		
	9-Aug-19	200	54900	0		
South Steyne Beach	6-Sep-19	110	53900	0		
Beach	23-Dec-19	41	54900	0		
	10-Jan-20	120	54400	0		
	16-Oct-19	120	54100	0.4	66098	Rose Bay (Royal Sydney Golf Club)
Tamarama Beach	24-Jan-20	39	54200	0		
	4-May-20	51	52100	0		
	1-Feb-20	41	53900	0	68108	Woonona (Popes Rd)
Thirroul Beach	1-Apr-20	48	53600	0.8	00000	Bellambi AWS
	7-Apr-20	41	54000	0	68228	
T : " 5 .	9-Aug-19	200	54300	0	66126	Collaroy (Long Reef Golf Club)
Turimetta Beach	23-Aug-19	39	54400	0		
Warriewood Beach	23-Aug-19	60	54500	0	66141	Mona Vale Golf Club
	6-Sep-19	95	54100	0.6		
Watsons Bay	24-Jan-20	58	53500	0	66099	Rose Bay (Royal Sydney Golf Club)
Whale Beach	6-Sep-19	53	53800	0	66128	Palm Beach (Sunrise Road)
	18-Jun-20	40	55800	1.2		
Woodford Bay	7-Aug-19	45	53100	0	66034	Abbotsford (Blackwall Point Rd)
Yarra Bay	28-Jan-20	76	53000	0.2	66037	Sydney Airport AMO






Appendix I : Chlorophyll-a trends at estuarine sites


Port Jackson






















00

Botany Bay


















Port Hacking








Appendix J : Water quality trends in lagoons

















Financial year



















































Appendix K: Intertidal communities

- Shoreline outfalls

Monitoring of rocky-intertidal communities under the shoreline outfall program assesses the potential ecological impact from the Shellharbour WWTP which discharges to the nearshore ocean environment. The structures of natural communities (without anthropogenic impacts) from two control sites were used in assessment of the Shellharbour outfall (impact) site (Volume 1 Figure 2-16). The Shellharbour outfall site is situated about 2 km north of the two control sites. The control sites were situated about 400 m apart. Shellharbour was the only WWTP that was measured under this program, as health and safety risks precluded access to intertidal rock platforms near the four other nearshore ocean outfall locations.

Rocky-intertidal communities are comprised of macro algae and macro invertebrate animals. These organisms will also colonise a variety of anthropogenic structures such as breakwaters, jetties, docks, groynes, dykes and seawalls (Crowe et al. 2000). Wave exposure is known to influence distribution and abundance of rocky-intertidal communities between exposed headlands and sheltered bays or inlets (Crowe et al. 2000). To control this natural influence, sites were selected that had similar levels of wave exposure. Rocky-intertidal community structure was monitored from wave-exposed ocean headland locations on naturally occurring rock platforms that could be safely accessed at low tide.

At each site, community composition and enumeration were recorded yearly during the period of late winter to late spring. Monitoring in this period reduces the influence of annual recruitment of most species of settling larvae that mainly happens in summer to autumn. Photographs of a 0.25 m² quadrat were taken within two hours either side of low tide. To help encapsulate variation between sites and across years, 14 randomly selected 0.25 m² quadrats were photographed between the low and high tide marks in the mid-littoral zone at each site visit. Using these photographs, counts were recorded for macro invertebrate taxa and estimates of percentage cover were made for macro algae. The taxonomic level recorded was based on morphological characteristics that could be seen with the naked eye and the level recorded is shown in the SIMPER 2019 output (Table K-2). Identification of macro invertebrate taxa and macro algae was checked against taxonomic works of Edgar (1997) and Dakin (1987).

Shoreline outfall discharges with documented measurable impacts on intertidal community structure are typically limited in spatial extent from 100 to 300 m (Fairweather 1990 and AWT 1998). These intertidal community structures were dominated by extensive covers of green macro algae. A pictorial example of a localised spatial impact of about 50 m² (Figure K-1) was formerly seen at Barrack Point outfall in 2001. At that time, an extensive cover of green macro algae occurred with few invertebrates (EP Consulting 2003). This was prior to upgrade works conducted at the Shellharbour WWTP in the early to mid-2000's.

The Bray-Curtis resemblance measure is focused on compositional changes in taxa identities (Anderson and Walsh 2013). As such, this is an appropriate choice since we understand the former measurable impact from near shore wastewater discharge at Shellharbour did cause a change in the composition of the intertidal rock platform community.

The PERMANOVA routine is designed to test whether it is reasonable to consider the existence of pre-defined groups given overall variability (Anderson et al. 2008).

An asymmetrical permutational multivariate analysis of variance test (PERMANOVA) was conducted with 'Control and Impact' locations treated as a fixed factor. 'Sites' were nested within 'Control and Impact', with 'Sites' treated as a random factor. The outfall site was the only site under the Impact location and the two sites were under the Control locations. A fourth root transformation was applied to the data prior to a Bray-Curtis dissimilarity matrix being constructed. This matrix was the basis for PERMANOVA testing with 9999 permutations run under a reduced model, with conservative Type III sums of squares inspected to base hypothesis decisions upon.

Asymmetrical PERMANOVA indicated there was no significant difference between 'Control and Impact' locations for the 2019 survey (Table K-3).

SIMPER analysis reflected a community structure dominated by invertebrates with a lesser contribution of macro algae at all three locations including the outfall location (Table K-4). The picture of the outfall site in 2019 (Figure K-1) reflects invertebrate dominance seen in SIMPER results, which is different to the green algal dominance recorded in 2001 prior to WWTP upgrade works (Figure K-2).

In summary the multivariate analyses of community structure of 2019 morphologically based intertidal rock platform community data suggested there was no measurable impact in the intertidal rock platform community near the outfall at Barrack Point from wastewater discharges from the Shellharbour WWTP. This outcome was supported by the differences apparent in the pictorial comparisons of 2001 and 2019. Context of 2019 data to the broader data collected back to 2008 is provided under the 2008 to 2019 data analysis below.

Table K-1 Asymmetrical PERMANOVA of 2019 intertidal assemblages

Permutational MANOVA

Sums of squares type: Type III (partial)
Fixed effects sum to zero for mixed terms

Permutation method: Permutation of residuals under a reduced model

Number of permutations: 9999

Factors

Name	Type	Levels
Control / Impact	Fixed	2
Site	Random	3

PERMANOVA table of results

						Unique	
Source	df	SS	MS	Pseudo-F	P(perm)	perms	P(MC)
Control / Impact	1	13024	13024	2.4238	0.3349	3	0.2108
Site(Control / Impact)	1	5373.5	5373.5	5.7498	0.0028	9958	0.0019
Res	39	36447	934.54				
Total	41	54845					

Estimates of components of variation

Source	Estimate	Sq.root
S(Control / Impact)	409.87	20.245
V(Site(Control / Impact))	317.07	17.806
V(Res)	934.54	30.57





Table K-2 SIMPER 2019 - intertidal assemblages by site

Control site-1 – 2019 Average sample similarity: 59%

Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%
Nerite (Nertidae <i>Nerita</i>)	2.18	20.32	4.48	34.34	34.34
Conniwinks (Lottorinidae Bembicium)	1.78	17.68	5.24	29.86	64.20
Periwinkles (Littorinidae Nodilitorina)	1.42	8.33	0.74	14.07	78.27
Brown algae (Phaeophyta)	1.33	4.81	0.48	8.12	86.39
False limpets & rock limpets (Patellogastropoda)	0.72	3.07	0.51	5.18	91.57
Zebra top shell (Trochidae Austrocochlea)	0.70	2.62	0.53	4.42	95.99
Barnacles (Cirripedia)	0.97	2.37	0.41	4.01	100.00

Control site-2 – 2019 Average sample similarity: 52%

Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%
Barnacles (Cirripedia)	4.69	19.78	1.53	38.38	38.38
Periwinkles (Littorinidae Nodilitorina)	1.90	8.08	0.67	15.69	54.07
Nerite (Nertidae Nerita)	1.58	7.19	1.15	13.95	68.02
Conniwinks (Lottorinidae Bembicium)	1.51	6.25	0.95	12.13	80.15
Brown algae (Phaeophyta)	1.87	6.12	0.70	11.89	92.03
Zebra top shell (Trochidae Austrocochlea)	0.81	1.91	0.44	3.71	95.74
False limpets & rock limpets (Patellogastropoda)	0.66	1.77	0.54	3.43	99.17
Oyster Borer (Muricidae Morula marginalba)	0.34	0.43	0.26	0.83	100.00

Outfall site – 2019 Average sample similarity: 59%

Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%
Conniwinks (Lottorinidae Bembicium)	2.40	15.38	4.02	26.18	26.18
False limpets & rock limpets (Patellogastropoda)	1.90	10.77	3.09	18.33	44.52
Zebra top shell (Trochidae Austrocochlea)	2.18	8.70	1.09	14.81	59.33
Red Algae (Rhodophyta)	2.28	7.40	0.78	12.60	71.93
Barnacles (Cirripedia)	1.40	4.41	0.74	7.51	79.44
Oyster Borer (Muricidae Morula marginalba)	0.88	3.74	0.95	6.36	85.81
Periwinkles (Littorinidae Nodilitorina)	1.11	3.22	0.53	5.48	91.29
Green Algae (Chlorophyta)	1.40	3.05	0.54	5.19	96.48
Brown algae (Phaeophyta)	0.86	1.32	0.33	2.26	98.74
Nerite (Nertidae Nerita)	0.41	0.74	0.35	1.26	100.00



Figure K-1 Barrack Point with a healthy intertidal rock platform community in 2019



Figure K-2 Barrack Point (in 2001) with an unhealthy intertidal rock platform community impacted by wastewater discharges from the Shellharbour WWTP prior to upgrade in the early to mid 2000's





Intertidal communities Shellharbour 2008 to 2019

Inclusion of yearly replicate samples from 2008 to 2019 allowed the factor 'Time' to be included in the above asymmetrical permutational analysis of variance test (PERMANOVA). Time was comprised of 2008, 2009, 2010, 2011, 2012, 2013, 2014, 2015, 2016, 2017, 2018 and 2019 surveys, which were conducted at varying times through late winter to late spring.

Asymmetrical PERMANOVA indicated there was no significant difference between 'Control and Impact' locations for the 2008 to 2019 period (Table K-3). However, differences between sites through time were indicated as significant results were returned for the 'Site (Control and Impact)' and 'Site (Control / Impact) x Time' factors (Table K-3).

The non-metric multidimensional scaling (nMDS) ordination routine of PRIMER was used to produce two and three dimensional ordination plots. In these plots the relative distance between samples is proportional to the relative similarity in taxonomic composition and abundance – the closer the points on the graph the more similar the community (Clarke 1993). That is, site samples with similar taxa lay closer together and site samples with a differing taxon composition lie farther apart. An unconstrained ordination procedure such as nMDS inevitably introduces distortion when trying to simultaneously represent the similarities between large numbers of samples in a few dimensions. The success of the procedure is measured by a stress value, which indicates the degree of distortion imposed. In the PRIMER software package a stress value of below 0.2 indicates an acceptable representation of the original data, although lower values are desirable. Where stress values are just above 0.2, the patterns displayed should be confirmed with other techniques such as PERMANOVA. The returned two-dimensional stress value was 0.22 and an improved lower stress value of 0.15 was observed for the three-dimensional ordination plot.

To understand the context of 2019 site data to that of previous years (2008 to 2018), site sample data were colour coded as shown in Figure K-3. Data patterns displayed in this two-dimensional nMDS ordination plot indicated 2008 to 2018 Control site-1 samples overlapped with 2008 to 2018 outfall site samples. The 2019 outfall samples also overlaid this mass of samples. While the Control site-1 samples from 2019 overlaid the agglomeration of Control site-1 2008 to 2018 samples. The 2019 Control-site 2 samples tended to spread from the edge of the agglomeration of Control site-2 2008 to 2019 indicating greater taxonomic variation to previous years. A three-dimensional nMDS ordination plot is presented in Figure K-4. It indicates a tighter agglomeration of samples than displayed in the two dimensional nMDS ordination plot in Figure K-3.

Under the nMDS routine, due to rank ordering of dissimilarities some detail can be hidden. This detail may be seen using a Principal Coordinates Analysis (PCO) routine as PCO is based upon original dissimilarities being projected onto axes in the space of the chosen resemblance measure (Anderson et al. 2008). As a check for any additional dimensionality in the multivariate data cloud, a PCO ordination plot was raised based on a fourth root transformation of the data and a Bray-Curtis resemblance measure. No additional dimensionality was indicated as the patterns between nMDS (Figure K-3, Figure K-4) and PCO ordination (Figure K-5) plots were very similar.

A Canonical Analysis of Principal Coordinates (CAP) ordination plot was also produced (Figure K-6). The CAP routine is designed to ask, 'Are there axes in the multivariate space that separate groups?' CAP is designed to purposely seek out and find groups even if differences occur in obscure directions and may not have been apparent from nMDS or PCO plots that provide views of the multivariate data cloud as a whole (Anderson et al. 2008). A similar pattern to that in the

nMDS (Figure K-3) and PCO (Figure K-5) ordination plots was displayed. This also suggested no hidden dimensionality, with good agreement between the nMDS, PCO and CAP ordination plots.

An additional run of the CAP routine was undertaken with placement of 2019 outfall samples onto the canonical axes of the existing CAP model from the initial run. Output from the second run indicated 2019 outfall samples were most similar to either the Outfall 2008 to 2018 samples or Control site 1 2008 to 2019 samples (Figure K-6). This result also reflected patterns displayed in the nMDS and PCO ordination plots (Figure K-4, Figure K-5 and Figure K-6).

The trend of taxonomic differences between sites situated close together on shorelines is known to occur and accounts for the differences between Control site-1 that is only 400 m from Control site-2 on the shoreline. It is mentioned by Underwood and Chapman (1998) who cite Underwood (1981) who states 'on exposed shores in New South Wales there are great differences in patterns of occupancy of space from one place to another not many metres away, even though these are not a function of gradients in wave action.'

In summary, a relatively stable equilibrium in rocky-intertidal community structure was indicated from these assessments of the 2008 to 2019 monitoring data at the three sites studied under the STSIMP. These results also suggest over the 2008 to 2019 period, no measurable impact had developed in the intertidal rock platform community near the outfall at Barrack Point from wastewater discharges from the Shellharbour WWTP as the community assemblage of the outfall site was very similar to Control site-1 for the 2008 to 2018 period. Results from Control site-2 represent natural variation in rocky-intertidal community structure that has been demonstrated to occur for closely spaced shoreline sites.








Figure K-3 Two-dimensional nMDS ordination plot of 2008 to 2019 intertidal rock platform community data




Figure K-4 Three-dimensional nMDS ordination plot of 2008 to 2019 intertidal rock platform community data








Figure K-5 PCO ordination plot of 2008 to 2019 intertidal rock platform community data - dimensional




Figure K-6 CAP ordination plot of intertidal rock platform community data (2008 to 2018 for Control site 1 and Control site 2 and 2008 to 2018 outfall site) with 2019 outfall samples (orange squares) predicted





Table K-3 Asymmetrical PERMANOVA of 2008 to 2019 intertidal assemblages

Sums of squares type: Type III (partial) Fixed effects sum to zero for mixed terms

Permutation method: Permutation of residuals under a reduced model

Number of permutations: 9999

Factors

Name	Type	Levels
Control / Impact	Fixed	2
Time	Fixed	12
Site	Random	3

PERMANOVA table of results

						Unique	
Source	df	SS	MS	Pseudo-F	P(perm)	perms	P(MC)
Control / Impact	1	54135	54135	0.72584	0.6671	3	0.6036
Time	11	67867	6169.7	1.2567	0.1999	9911	0.2459
Site(Control / Impact)	1	73275	73275	101.58	0.0001	9950	0.0001
Control / ImpactxTime	11	48254	4386.7	0.89421	0.6668	9901	0.6372
TimexSite(Control / Impact)	11	53227	4838.8	6.7082	0.0001	9897	0.0001
Res	466	3.3614E+05	721.33				
Total	E04	C 2040E . 0E					

Total 501 6.3919E+05

Estimates of components of variation

Source	Estimate	Sq.root
S(Control / Impact)	-91.177	-9.5486
S(Time)	33.744	5.8089
V(Site(Control / Impact))	439.06	20.954
S(Control / ImpactxTime)	-27.814	-5.2739
V(TimexSite(Control / Impact))	298.31	17.272

V(Res) 721.33 26.858





Table K-4 CAP analysis of 2008 to 2018 intertidal assemblages with 2019 outfall site samples predicted

Factor level for new samples group: Outfall-2019

Number of samples: 488

Choice of m: 3

CANONICAL ANALYSIS

Correlations

Eigenvalue Correlation Corr.Sq.
1 0.8122 0.6597
2 0.25 0.0625

DIAGNOSTICS

m prop.G ssres d_1^2 d_2^2 %correct 3 0.7144 1.2945 0.6597 0.0625 65.984

Cross Validation

Leave-one-out Allocation of Observations to Groups (for the choice of m: 3)

Classified

Orig. group	Control 1-2008 to	Control 2-2008 to	Outfall-2008 to	Total	%correct
	2019	2019	2018		
Control 1-2008 to 2019	75	18	75	168	44.643
Control 2-2008 to 2019	13	151	0	164	92.073
Outfall-2008 to 2018	54	6	96	156	61.538

Total correct: 322/488 (65.984%) Mis-classification error: 34.016%

Individual samples that were mis-classified

Sample	Orig.group	Class.group
Control 1-2018-1	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2018-4	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2018-5	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2018-6	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2018-10	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2018-11	Control 1-2008 to 2019	Control 2-2008 to 2019
Control 1-2018-12	Control 1-2008 to 2019	Control 2-2008 to 2019
Control 1-2018-13	Control 1-2008 to 2019	Control 2-2008 to 2019
Control 1-2018-14	Control 1-2008 to 2019	Control 2-2008 to 2019
Control 1-2008-1	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2008-2	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2008-3	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2008-4	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2008-5	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2008-6	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2008-7	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2008-8	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2008-9	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2008-10	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2008-11	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2008-12	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2009-2	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2009-4	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2009-5	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2009-7	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2009-9	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2009-10	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2009-12	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2009-13	Control 1-2008 to 2019	Outfall-2008 to 2018





Control 1-2009-14	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2010-1	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2010-2	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2010-4	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2010-5	Control 1-2008 to 2019	Control 2-2008 to 2019
Control 1-2010-6	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2010-10	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2010-11	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2010-12	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2011-3	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2011-4	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2011-5	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2011-7	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2011-9	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2011-10	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2011-12	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2012-1	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2012-2	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2012-6	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2012-7	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2012-9	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2012-11	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2012-14	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2013-7	Control 1-2008 to 2019	Control 2-2008 to 2019
Control 1-2013-9	Control 1-2008 to 2019	Control 2-2008 to 2019
Control 1-2013-13	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2014-1	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2014-2	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2014-3	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2014-4	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2014-5	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2014-7	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2014-8	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2014-10	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2014-11	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2014-12	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2014-13	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2014-14	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2015-3	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2015-5	Control 1-2008 to 2019	Control 2-2008 to 2019
Control 1-2015-8	Control 1-2008 to 2019	Control 2-2008 to 2019
Control 1-2015-10	Control 1-2008 to 2019	Control 2-2008 to 2019
Control 1-2015-11	Control 1-2008 to 2019	Control 2-2008 to 2019
Control 1-2015-14	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2016-1	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2016-2	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2016-5	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2016-12	Control 1-2008 to 2019	Control 2-2008 to 2019
Control 1-2017-1	Control 1-2008 to 2019	Control 2-2008 to 2019
Control 1-2017-2	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2017-5	Control 1-2008 to 2019	Control 2-2008 to 2019
Control 1-2017-6	Control 1-2008 to 2019	Outfall 2008 to 2018
Control 1-2017-7	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2017-8	Control 1-2008 to 2019	Control 2-2008 to 2019
Control 1-2017-9	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2017-14	Control 1-2008 to 2019	Control 2-2008 to 2019
Control 1-2019-2	Control 1-2008 to 2019	Outfall 2008 to 2018
Control 1 -2019-5	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2019-6	Control 1-2008 to 2019	Control 2-2008 to 2019





Control 1-2019-7	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2019-8	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2019-9	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2019-10	Control 1-2008 to 2019	Outfall-2008 to 2018
Control 1-2019-11	Control 1-2008 to 2019	Control 2-2008 to 2019
Control 2-2008-2	Control 2-2008 to 2019	Control 1-2008 to 2019
Control 2-2008-6	Control 2-2008 to 2019	Control 1-2008 to 2019
Control 2-2008-10	Control 2-2008 to 2019	Control 1-2008 to 2019
Control 2-2008-13	Control 2-2008 to 2019	Control 1-2008 to 2019
Control 2-2008-14	Control 2-2008 to 2019	Control 1-2008 to 2019
Control 2-2019-1	Control 2-2008 to 2019	Control 1-2008 to 2019
Control 2-2019-2	Control 2-2008 to 2019	Control 1-2008 to 2019
Control 2-2019-3	Control 2-2008 to 2019	Control 1-2008 to 2019
Control 2-2019-4	Control 2-2008 to 2019	Control 1-2008 to 2019
Control 2-2019-5	Control 2-2008 to 2019	Control 1-2008 to 2019
Control 2-2019-7	Control 2-2008 to 2019	Control 1-2008 to 2019
Control 2-2019-8	Control 2-2008 to 2019	Control 1-2008 to 2019
Control 2-2019-9	Control 2-2008 to 2019	Control 1-2008 to 2019
Outfall-2018-1	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2018-2	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2018-3	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2018-5	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2018-6	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2018-7	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2018-9	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2018-11	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2018-12	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2018-13	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2018-14	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2008-12	Outfall 2008 to 2018	Control 1-2008 to 2019
Outfall-2010-7	Outfall 2008 to 2018	Control 1-2008 to 2019
Outfall-2011-1	Outfall-2008 to 2018	Control 2-2008 to 2019
Outfall-2011-2	Outfall 2008 to 2018	Control 2-2008 to 2019
Outfall-2011-3	Outfall 2008 to 2018	Control 1-2008 to 2019
Outfall-2011-7	Outfall 2008 to 2018	Control 2-2008 to 2019
Outfall-2011-9	Outfall 2008 to 2018	Control 1-2008 to 2019
Outfall-2011-10 Outfall-2011-11	Outfall-2008 to 2018 Outfall-2008 to 2018	Control 1-2008 to 2019 Control 1-2008 to 2019
Outfall-2011-13	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2012-2	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2012-4	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2012-4	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2013-3	Outfall-2008 to 2018	Control 2-2008 to 2019
Outfall-2013-12	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2014-1	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2014-2	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2014-5	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2014-8	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2014-9	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2014-10	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2015-1	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2015-4	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2015-6	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2015-8	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2015-13	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2015-14	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2016-1	Outfall-2008 to 2018	Control 2-2008 to 2019
Outfall-2016-3	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2016-4	Outfall-2008 to 2018	Control 1-2008 to 2019





Outfall-2016-5	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2016-6	Outfall-2008 to 2018	Control 2-2008 to 2019
Outfall-2016-9	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2016-11	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2016-12	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2016-13	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2016-14	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2017-1	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2017-2	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2017-3	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2017-4	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2017-5	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2017-6	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2017-8	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2017-9	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2017-10	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2017-11	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2017-12	Outfall-2008 to 2018	Control 1-2008 to 2019
Outfall-2017-14	Outfall-2008 to 2018	Control 1-2008 to 2019

PERMUTATION TEST

trace statistic = (tr(Q_m'HQ_m))

first squared canonical correlation = (delta_1^2)

tr(Q_m'HQ_m): 0.72222 P: 0.0001 delta_1^2: 0.65974 P: 0.0001 No. of permutations used: 9999

NEW SAMPLES

Canonical coordinate scores for New Sam		w Samples
Sample	CAP1	CAP2
Outfall-2019-1	-0.0047	-0.0049
Outfall-2019-2	0.0379	-0.0053
Outfall-2019-3	0.0403	-0.009
Outfall-2019-4	0.0088	0.0005
Outfall-2019-5	0.0111	-0.0095
Outfall-2019-6	-0.0105	0.0009
Outfall-2019-7	-0.0195	-0.0086
Outfall-2019-8	0.0283	0.0005
Outfall-2019-9	0.0353	-0.0068
Outfall-2019-10	0.0238	-0.0059
Outfall-2019-11	0.046	-0.0034
Outfall-2019-12	0.0126	-0.0073
Outfall-2019-13	0.0366	0.0047
Outfall-2019-14	0.0402	0.0046

New sample classification

Sample	Group
Outfall-2019-1	Control 1-2008 to 2019
Outfall-2019-2	Outfall-2008 to 2018
Outfall-2019-3	Outfall-2008 to 2018
Outfall-2019-4	Control 1-2008 to 2019
Outfall-2019-5	Control 1-2008 to 2019
Outfall-2019-6	Control 1-2008 to 2019
Outfall-2019-7	Control 2-2008 to 2019
Outfall-2019-8	Outfall-2008 to 2018
Outfall-2019-9	Outfall-2008 to 2018
Outfall-2019-10	Control 1-2008 to 2019
Outfall-2019-11	Outfall-2008 to 2018



Outfall-2019-12 Control 1-2008 to 2019
Outfall-2019-13 Outfall-2008 to 2018
Outfall-2019-14 Outfall-2008 to 2018





Intertidal rock platform communities


As a check of potential change in community structure of intertidal rock platforms at test sites, a comparison was made between control sites and other sites situated below urban catchments. This check was conducted under Principal Coordinates Analysis (PCO). PCO is an ordination technique that projects points onto axes that minimise the residual variation in the space of a chosen dissimilarity measure (Anderson et al. 2008). The user chooses the number of axes to be included in the output, but usually the first 2 or 3 axes contain most of the percent variation explained. In the analysis presented here, PCO was based on a Bray-Curtis distance measure matrix. The Bray-Curtis resemblance measure is focused on compositional changes in taxa identities (Anderson and Walsh 2013). The choice of this measure is considered appropriate as a change in taxonomic composition was recorded after remediation of the wastewater system (Sydney Water 2012). A separate analysis was conducted for each salinity zone.


The PCO output allowed control chart style visualisation of centroids in Bray-Curtis space for each site by plotting output for PCO axis 1 against year. This explained about 65% of the variation for the low salinity zone and about 27% of the variation for the high salinity zone. This indicated the low salinity analysis described more variation in data of low salinity sites (86% of variation explained by the first two PCO axes) compared with that for the high salinity sites (47% of variation explained by the first two PCO axes) (Figure L-1 and Figure L-2) over the 1998 to 2019 period.


Test sites within higher salinity zones in 2019 were grouped near or within the range of variation recorded for higher salinity control sites. However, those in the lower salinity zone were well separated, particularly in the Parramatta River and Cooks River, from the recorded range of variation for the lower salinity control sites. In contrast, test sites within the Georges River (Como Woronora River GR15; Kyle Bay Georges River GR115; Edith Bay Georges River GR175), with the exception of Salt Pan Creek GR18, showed consistent improvement in the last three years and now sit within the range of control sites. This suggests the 2019 community structure in the lower salinity zone at most sites (Kissing Point Bay Parramatta River PJ025; Silverwater Bridge Parramatta River PJ01; Woolwich Baths Lane Cove River PJ05; Salt Pan Creek Georges River GR18; Alexandra Canal Cooks River CR04; Wolli Creek Cooks River CR06) were impaired. The exception being an improvement in the trend for the Hawthorn Canal arm within Iron Cove (PJ082).












Black line colour represents control sites: other line colours represent test sites

Figure L-1 Relatively lower salinity zone with year plotted against Principal Coordinates
Analysis axis 1 of distance among centroids for sites







Black line colour represents control sites: other line colours represent test sites

Figure L-2 Relatively higher salinity zone with year plotted against Principal Coordinates Analysis axis 1 of distance among centroids for sites



00

Settlement panels

Settlement panels were used to supplement intertidal rock platform measurements and provide a focus on colonisation of intertidal larvae from the swimming juvenile life stages. Settlement panels were deployed at a number of sites that each included a large muddy intertidal area with mangroves. These areas of the estuaries did not have regular wave activity. The settlement panels consisted of weathered hardwood fence palings (weathered to remove tannins) that were vertically hammered into the mud at an intertidal height just below the lowest growing mangroves, and were left for four months to allow intertidal organisms to settle. After that time, they were removed and measured for the area covered by barnacles. Panels were deployed twice a year, during late spring and late autumn.

Barnacles were the dominant animal that settled and were a mixture of small types like *Elminius* and *Chamaesipho*, as well as some larger animals like *Balanus*. The relatively short deployment time of approximately four months, was inadequate for taxa such as snails (Mollusca) to develop to a sufficient size compared to barnacles that developed in a relatively shorter time where conditions were suitable for barnacle settlement. Previous analysis by Sydney Water (2012) showed reductions in barnacle cover (for example Rushcutters Bay PJ33) following sewer remediation suggesting higher levels of barnacle cover to be a possible indicator of wastewater overflows in wave-sheltered areas of the estuaries around Sydney.

In wave exposed areas of the coast and outer estuaries where there is regular wave occurrence, barnacles naturally grow on hard substrates and are not an indicator of the presence of wastewater. An estimate of barnacle cover was formed by multiplying the average size of barnacles with measured abundance.

Analysis of variance (ANOVA) indicated significant differences between sites situated in relatively higher salinity waters of the outer estuaries of Sydney (df = 9, MS = 684.410, F = 16.73, p <0.0001). A multiple mean (SNK) comparison indicated two sites on the Georges River being significantly different from each other and the remaining overlapping group of sites Figure L-3.

Site	Estimate
GR01	35.3131
GR085	17.4341
PJ33	9.8594
PJ295	0.8831
PHE05	0.3500
PJ13	0.1155
PJ28	0.05613
PJ315	0.05000
PW10	0.04363
PH05	0.01071

Figure L-3 Multiple mean comparison groupings of relatively high salinity locations for 2019-20. (means covered by the same bar are not significantly different)





ANOVA indicated significant differences between sites situated in relatively lower salinity waters of the inner estuaries of Sydney (df = 12, MS = 3540.468, F = 22.51, p <0.0001). A multiple mean (SNK) comparison test indicated one significantly different site on the Georges River from the remaining four overlapping groups of sites (Figure L-4).

Site	Estimate
GR15	68.7618
PJ082	43.2218
GR175	33.9784
PJ05	23.8369
GR115	19.8739
PJ01	7.9947
CR06	2.8481
PJ025	2.2256
CR04	1.0125
GR18	0.2619
PH10	0.09063
NCC02	0.01875
NCC01	0.006250

Figure L-4 Multiple mean comparison groupings of relatively lower salinity locations for 2019-20. (means covered by the same bar are not significantly different)



Appendix M : Hawkesbury-Nepean River water quality trends

The Hawkesbury-Nepean River sites are presented in the following order from upstream to downstream:

- N92: Nepean River at Maldon Weir
- N75: Nepean River at Sharpes Weir
- N67: Nepean River at Wallacia Bridge
- N57: Nepean River at Penrith Weir
- N51: Nepean River opposite Fitzgerald Creek
- N48A: Nepean River at Smith Road
- N44: Nepean River at Yarramundi Bridge
- N42: Hawkesbury River at North Richmond
- N39: Hawkesbury River at Freemans Reach
- NS04A: Lower South Creek at Fitzroy Bridge
- N35: Hawkesbury River at Wilberforce
- NC11A: Lower Cattai Creek at Cattai Ridge Road
- N3001: Hawkesbury River off Cattai SRA
- N26: Hawkesbury River at Sackville Ferry
- N2202: Lower Colo River at Putty Road
- N18: Hawkesbury River at Leets Vale
- NB13: Berowra Creek at Calabash Bay
- NB11: Berowra Creek off Square Bay

The water quality plots are presented in the following groups and order of analytes:

Nutrients, chlorophyll-a and algae

- Ammonia nitrogen
- Oxidised nitrogen
- Total nitrogen
- Filterable total phosphorus
- Total phosphorus
- Chlorophyll-a
- Total algal biovolume
- Blue-green algal biovolume
- Toxic blue-green algal biovolume
- Toxic blue-green algal count


Other physico-chemical analytes

- Conductivity
- Dissolved oxygen (mg/L)
- Dissolved oxygen saturation (%)
- pH
- Temperature
- Turbidity






N92: Nepean River at Maldon Weir




The ordination plot from Principal Components Analysis suggested water quality of the Nepean River at Maldon Weir in 2019-20 was similar in variability to that recorded in the period 2010 to 2019.


Nutrients, chlorophyll-a and algae








































































N75: Nepean River at Sharpes Weir





The ordination plot from Principal Components Analysis suggested water quality of the Nepean River at Sharpes Weir in 2019-20 was similar to that recorded in the period 2010 to 2019, although these more recent samples appeared to have relatively higher concentrations of nitrogen parameters.













































Other physico-chemical analytes


















N67: Nepean River at Wallacia Bridge





The ordination plot from Principal Components Analysis suggested water quality of the Nepean River at Wallacia Bridge in 2019-20 was similar to that recorded in the period 2010 to 2019.













































Other physico-chemical analytes


























N57: Nepean River at Penrith Weir





The ordination plot from Principal Components Analysis suggested water quality of the Nepean River at Penrith Weir in 2019-20 was similar to that recorded in the period 2010 to 2019.








































































N51: Nepean River opposite Fitzgerald Creek





The ordination plot from Principal Components Analysis suggested water quality of the Nepean River opposite Fitzgeralds Creek in 2019-20 was similar to that recorded in the period 2010 to 2019.








































































N48A: Nepean River at Smith Road





The ordination plot from Principal Components Analysis suggested water quality of the Nepean River at Smith Road in 2019-20 was similar to that recorded in the period 2010 to 2019.












































Other physico-chemical analytes





















N44: Nepean River at Yarramundi Bridge





The ordination plot from Principal Components Analysis suggested water quality of the Nepean River at Yarramundi Bridge in 2019-20 was similar to that recorded in the period 2010 to 2019.






































































N42: Hawkesbury River at North Richmond





The ordination plot from Principal Components Analysis suggested water quality of the Hawkesbury River at North Richmond in 2019-20 was similar to that recorded in the period 2010 to 2019.






































































N39: Hawkesbury River at Freemans Reach





The ordination plot from Principal Components Analysis suggested water quality of the Hawkesbury River at Freemans Reach in 2019-20 was similar to that recorded in the period 2010 to 2010.






































































NS04A: Lower South Creek at Fitzroy Bridge





The ordination plot from Principal Components Analysis suggested water quality of Lower South Creek at Fitzroy Bridge in 2019-20 had relatively similar variability to that recorded in the period 2010 to 2019.








































































N35: Hawkesbury River at Wilberforce





The ordination plot from Principal Components Analysis suggested water quality of the Hawkesbury River at Wilberforce in 2019-20 was similar to that recorded in the period 2010 to 2019.






































































NC11A: Lower Cattai Creek at Cattai Ridge Road





The ordination plot from Principal Components Analysis suggested water quality of the Lower Cattai Creek at Cattai Ridge Road in 2019-20 had similar variability to that recorded in the period 2010 to 2019.








































































N3001: Hawkesbury River off Cattai SRA





The ordination plot from Principal Components Analysis suggested water quality of the Hawkesbury River off Cattai SRA in 2019-20 was similar to that recorded in the period 2010 to 2019.








































































N26: Hawkesbury River at Sackville Ferry





The ordination plot from Principal Components Analysis suggested water quality of the Hawkesbury River at Sackville ferry in 2019-20 was similar to that recorded in the period 2010 to 2019.
































































N2202: Lower Colo River at Putty Road





The ordination plot from Principal Components Analysis suggested water quality of the Lower Colo River at Putty Road in 2019-20 was generally similar to that recorded in the period 2010 to 2019.








































































N18: Hawkesbury River at Leets Vale





The ordination plot from Principal Components Analysis suggested water quality of the Hawkesbury River at Leets Vale in 2019-20 had similar variability to that recorded in the period 2010 to 2019.






































































NB13: Berowra Creek at Calabash Bay





The ordination plot from Principal Components Analysis suggested water quality of Berowra Creek at Calabash Bay in 2019-20 was similar to that recorded in the period 2010 to 2019.
































Note: Insufficient data to draw a plot on blue-green algal biovolume


Note: Insufficient data to draw a plot on toxic blue-green algal biovolume


Note: Insufficient data to draw a plot on toxic blue-green algal count

























NB11: Berowra Creek off Square Bay





The ordination plot from Principal Components Analysis suggested water quality of Berowra Creek off Square Bay in 2019-20 was generally similar to that recorded in the period 2010 to 2019.
































Note: Insufficient data to draw a plot on blue-green algal biovolume


Note: Insufficient data to draw a plot on toxic blue-green algal biovolume


Note: Insufficient data to draw a plot on toxic blue-green algal count

























Stream health in response to discharge from the inland WWTPs are presented in order of upstream to downstream sites:

- Picton WWTP
- West Camden WWTP
- Wallacia WWTP
- Penrith WWTP
- Winmalee WWTP
- North Richmond WWTP
- St Mary's WWTP
- Quakers Hill WWTP
- Riverstone WWTP
- Castle Hill WWTP
- Rouse Hill WWTP
- Hornsby Heights WWTP
- West Hornsby WWTP





Stream health near Picton WWTP

The SIGNAL-SG plot for the Nepean River provided an assessment of stream health. This plot was based on macroinvertebrate identification and counting results expressed as SIGNAL-SG scores and allows a visual comparison of data collected from 2019–20 against that collected between 1995 and 2019. This comparison suggests downstream stream health was maintained at a level comparable to that of the upstream site recorded over the 1995 to 2019 period indicating wastewater discharge from Picton WWTP did not have a measurable negative impact on stream health during 2019-20 (Figure N-1).

A comparison of the upstream-downstream SIGNAL-SG scores for 2019-20 samples under ttests returned non-significant test outcomes (Table N-1) and confirmed the visual trend for 2019-20.

As no measurable negative impact on downstream stream health was detected, no further data analysis was undertaken.




Figure N-1 Stream health of Nepean River near Picton WWTP

Table N-1 t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from the Nepean River near Picton WWTP

Method	Variances	DF	t Value	Pr > t
Pooled	Equal	14	1.38	0.1878

Equality of Variances					
Method	Num DF	Den DF	F Value	Pr > F	
Folded F	7	7	1.74	0.4830	





Stream health near West Camden WWTP

The SIGNAL-SG plots provided assessments of stream health for both Matahil Creek near West Camden WWTP and in the Nepean River upstream-downstream of the confluence with Matahil Creek. These plots were based upon macroinvertebrate identification and counting results expressed as SIGNAL-SG scores and allows a visual comparison of data collected from 2019–20 against that collected between 2004 to 2019 for the Matahil Creek and 1995 to 2019 for the Nepean River. These visual comparisons suggest downstream stream health was maintained in the Nepean River over 1995 to 2019 (Figure N-3, Table N-3). A localised impact in stream health was indicated for the Matahil Creek in 2019-20 (Figure N-2).

A comparison of the upstream-downstream SIGNAL-SG scores for 2019-20 samples under t-tests returned a significant test outcome for Matahil Creek (Table N-2) and a non-significant test outcome for the Nepean River, which confirmed the visual trends of respective SIGNAL-SG plots (Figure N-2 and Figure N-3).

As a measurable negative impact on downstream stream health was detected on Matahil Creek, further data analysis was undertaken.




Figure N-2 Stream health of Matahil Creek near West Camden WWTP

Table N-2 t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from Matahil Creek near West Camden WWTP

Method	Variances	DF	t Value	Pr > t
Pooled	Equal	7	-4.67	0.0023

Equality of Variances					
Method	Num DF	Den DF	F Value	Pr > F	
Folded F	5	2	4.02	0.4222	








Figure N-3 Stream health of the Nepean River near West Camden WWTP

Table N-3 t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from the Nepean River near West Camden WWTP

Method	Variances	DF	t Value	Pr > t
Pooled	Equal	10	0.53	0.6076

Equality of Variances					
Method Num DF Den DF F Value Pr > F					
Folded F	5	5	3.31	0.2154	

Edge habitat samples were collected consistently enough from Matahil Creek to allow multivariate analysis for the monitoring period 2004 to 2020. Distinct groups of samples separated by site were evident for Matahil Creek in the two-dimensional ordination plot (Figure N-4).

The ordination pattern was confirmed in the corresponding tree diagram (dendrogram) from classification analysis as the first division separated all upstream site samples from all downstream site samples (Figure N-5). This initial separation also occurred at a quite low similarity of 12% (Figure N-5) compared with all Nepean River sites samples which exhibited a greater initial similarity level of 37% (Figure N-8).

The clear separation of Matahil Creek sites was also evident in the corresponding shade plot (Figure N-6) where downstream samples displayed less diversity when compared to the upstream site. The shade plot displayed a few taxa in common between the two sites such as the freshwater snail Physidae *Physela*, the caddisfly larvae Leptoceridae *Triplectides* and the non-biting midge larvae Chironimidae *Chironomus*. The corresponding SIGNAL-SG grades showed that dominant taxa that occurred downstream have lower SIGNAL-SG grades than those of the





upstream site (Figure N-6), which is reflected in the separation of site SIGNAL scores displayed in Figure N-2.

The BVSTEP routine was used to find a subset of taxa whose multivariate pattern matched that of the full dataset with 17 taxa identified for the edge habitat (Table N-8) out of 134 taxa. These taxa reflected those taxa which formed the main patterns within the shade plot (Figure N-6).

The PERMDISP analysis indicated a similar pattern of dispersion (spacing between same site samples) for the two sites (Table N-6). This outcome suggests the variability in taxonomic makeup of samples collected over time was at similar levels for both sites through the period tested (2005 to 2020). This result then also implies subsequent results of ANOSIM tests are focused on community structure differences between sites.

The ANOSIM test run on the factor 'Site' returned a high range value (R = 0.998; P = 0.001) confirming community structure was distinct at each site (Table N-4).

To further explore community structure, hypothesis testing was conducted with PERMANOVA. The PERMANOVA model comprised the fixed factors 'Site' and 'Year'. 'Year' represented samples collected in years between 2005 and 2020. 'Site' had two levels, upstream and downstream. A statistically non-significant 'Site x Year' interaction was returned (Table N-5). This non-significant result allowed us to view the 'Site' and 'Year' results. A statistically non-significant 'Year' result was returned, however 'Site' resulted in a statistically significant result indicating differences exist between the upstream and downstream sites.

A second run of ANOSIM based on Site-period sample groups displayed in above ordination plots returned a significant global R-value at a high level of 0.772 (Table N-7). Pairwise tests indicated the four upstream versus downstream comparisons also had high level R-values (close to or equaling the maximum R-value of 1). In contrast, the same site comparison of the two time periods within each site were non-significant and returned a low level R-values. These pairwise test results suggest clear differences in assemblage structure between upstream and downstream sites, and that each site had a relatively stable community structure through time (Table N-7).

These results suggested downstream community structure in Matahil Creek was consistently altered by wastewater discharge from West Camden WWTP.








Figure N-4 Two-dimensional ordination plot of freshwater macroinvertebrate edge habitat community structure of Matahil Creek upstream and downstream sites of West Camden WWTP




Figure N-5 Tree diagram of freshwater macroinvertebrate edge habitat community structure of Matahil Creek upstream and downstream sites of West Camden WWTP





Tests for differences between unordered Site groups

Global Test

Sample statistic (R): 0.998

Significance level of sample statistic: 0.01%

Number of permutations: 9999 (Random sample from a large number)

Number of permuted statistics greater than or equal to R: 0

Table N-5 PERMANOVA test of 'Site' and 'Year' factors for edge habitat of Matahil Creek upstream and downstream sites of West Camden WWTP

Sums of squares type: Type III (partial) Fixed effects sum to zero for mixed terms

Permutation method: Permutation of residuals under a reduced model

Number of permutations: 9999

Factors

Name Type Levels Site Fixed 2 Year Fixed 16

PERMANOVA table of results

						Unique
Source	df	SS	MS	Pseudo-F	P(perm)	perms
Site	1	63017	63017	41.471	0.0001	9940
Year	15	26642	1776.1	1.1689	0.0625	9750
SitexYear	14	23248	1660.6	1.0928	0.1945	9712
Res	28	42547	1519.5			
Total 58	1.606	31E+05				

Estimates of components of variation

Source	Estimate	Sq.root
S(Site)	2254.9	47.486
S(Year)	70.241	8.381
S(SitexYear)	74.799	8.6487
V(Res)	1519.5	38.981

Table N-6 PERMDISP test of 'Site' factor for edge habitat of Matahil Creek upstream and downstream sites of West Camden WWTP

Group factor: Site

Number of permutations: 9999

Number of groups: 2 Number of samples: 59

DEVIATIONS FROM CENTROID

F: 0.31846 df1: 1 df2: 57

P(perm): 0.5806

MEANS AND STANDARD ERRORS

 Group
 Size
 Average
 SE

 Downstream
 31
 39.601
 1.2212

 Upstream
 28
 38.722
 0.92456





Table N-7 ANOSIM test of 'Site period' samples for edge habitat of Matahil Creek upstream and downstream sites of West Camden WWTP

Tests for differences between unordered Site period groups

Global Test

Sample statistic (R): 0.772

Significance level of sample statistic: 0.01%

Number of permutations: 9999 (Random sample from a large number)

Number of permuted statistics greater than or equal to R: 0

Pairwise Tests

1 dii 1100 1 000					
	R	Significance	Possible	Actual	Number >=
Groups	Statistic	Level %	Permutations	Permutations	Observed
Downstream 2005 to 2016, Downstream 2017 to 2020	-0.019	56.3	2629575	9999	5632
Downstream 2005 to 2016, Upstream 2005 to 2016	0.998	0.01	Very large	9999	0
Downstream 2005 to 2016, Upstream 2017 to 2020	0.99	0.01	20475	9999	0
Downstream 2017 to 2020, Upstream 2005 to 2016	1	0.01	2629575	9999	0
Downstream 2017 to 2020, Upstream 2017 to 2020	1	0.3	330	330	1
Upstream 2005 to 2016, Upstream 2017 to 2020	0.091	26.1	20475	9999	2604

Table N-8 Genera subset whose multivariate pattern matches full genera set of the edge habitat of Matahil Creek upstream and downstream sites of West Camden WWTP

Subset of 17 (correlation 0.951) genera from edge habitat whose pattern matches that of the full set of 134 genera identified with the same subset found on 9 runs from 50 random start runs. Each run was based on three randomly selected genera. Genera were:

Chironomidae *Chironomus*, Coenagrionidae *Ischnura*, Physidae *Physella*, Belostomatidae *Diplonychus*, Chironomidae *Cricotopus*, Chironomidae *Cricotopus*




Figure N-6 Shade plot of freshwater macroinvertebrate edge habitat community structure of Matahil Creek upstream and downstream sites of West Camden WWTP

At both upstream and downstream sites on the Nepean-River, edge habitat data was collected consistently enough through time (less sample collection gaps outlined in Volume 1 (Table 3-2) to allow multivariate analysis.

The Nepean River edge habitat data pattern was visually displayed in a three-dimensional nMDS ordination plot, as the two-dimensional plot had a poor (stress) value of 0.26. A stress value of that size potentially represents points being placed almost arbitrarily in two-dimensional space and suggests that there is no clear pattern of upstream downstream differences in the data. Addition of a third dimension provided a more acceptable stress value of 0.19. Data points were colour coded by Site-time periods (Figure N-7). Addition of a third dimension did not reveal a clear separation of groups of upstream and downstream samples in the corresponding ordination plot (Figure N-7).

The lack of a clear upstream downstream site pattern in the ordination plot was confirmed in the corresponding tree diagram (Figure N-8). Initial separation of samples occurred at a moderate level of similarity (37%) (Figure N-8).

The shade plot of the Nepean River edge habitat lacked a distinct site difference in the taxa pattern as seen for the Matahil Creek sites. Rather a less distinct difference between the 2018 to 2020 and 1995 to 2018 periods was apparent for both sites (Figure N-9). Looking at corresponding SIGNAL-SG grades revealed a mix of mid-range grades in both periods for both sites (Figure N-9).

The BVSTEP routine was used to find a subset of taxa whose multivariate pattern matched that of the full dataset with 66 taxa for the edge habitat (Table N-13) out of 178 taxa. This subset of taxa formed the main visual pattern in the respective shade plot (Figure N-9).

The PERMDISP analysis indicated a similar pattern of dispersion (spacing between same site samples) for the two sites (Table N-11). This suggests the variability in taxonomic make-up of samples collected over time was at similar levels for both sites through the period tested (1995 to 2020). This result then also implies subsequent results of ANOSIM tests are focused on community structure differences between sites rather than within.

An ANOSIM test was run on the factor 'Site'. The returned ANOSIM R-value was close to zero (0.108) (Table N-9), implying there was a lack of clearly different taxonomic assemblages present at each site, which was in contrast to the distinct community structure differences shown for Matahil Creek.

To further explore community structure, hypothesis testing was conducted using PERMANOVA (Table N-10). This model included the fixed factors 'Site' and 'Year'. 'Year' represented samples collected in years between 1995 and 2020 whereas 'Site' had two levels, upstream and downstream. A statistically non-significant 'Site x Year' interaction was returned (Table N-10). This non-significant result allowed us to view the 'Site' and 'Year' results. Significant results were returned for 'Site' and 'Year'. The components of variation output indicated 'Year' explained approximately eight times the variation than explained by 'Site' (Table N-10).

A second run of ANOSIM based on Site-period sample groups returned a significant global R-value at a low level of effectively zero (0.083) (Table N-12). Pairwise test outputs were non-significant for five of the six comparisons.

Both SIGNAL-SG and multivariate analysis results suggested downstream community structure in Matahil Creek was consistently altered by wastewater discharge from West Camden WWTP but this impact did not extend as far as the Nepean River.








Figure N-7 Dimensions 1 and 2 of three-dimensional ordination plot of freshwater macroinvertebrate edge habitat community structure of sites upstream-downstream of Nepean River at the confluence of Matahil Creek into which West Camden WWTP discharges




Figure N-8 Tree diagram of freshwater macroinvertebrate edge habitat community structure of sites upstream-downstream of Nepean River at the confluence of Matahil Creek into which West Camden WWTP discharges



Tests for differences between unordered Site groups

Global Test

Sample statistic (R): 0.108

Significance level of sample statistic: 0.02%

Number of permutations: 9999 (Random sample from a large number)

Number of permuted statistics greater than or equal to R: 1

Table N-10 PERMANOVA test of 'Site' and 'Year' factors for edge habitat of upstreamdownstream sites of Nepean River at the confluence of Matahil Creek into which West Camden WWTP discharges

Sums of squares type: Type III (partial) Fixed effects sum to zero for mixed terms

Permutation method: Permutation of residuals under a reduced model

Number of permutations: 9999

Factors

Name Type Levels Site Fixed 2 Year Fixed 26

PERMANOVA table of results

						Unique
Source	df	SS	MS	Pseudo-F	P(perm)	perms
Site	1	2858.2	2858.2	2.4597	0.0011	9910
Year	25	62030	2481.2	2.1353	0.0001	9618
SitexYear	19	20830	1096.3	0.94346	0.7645	9664
Res	38	44156	1162			
Total 83	1.312	21E+05				

Estimates of components of variation

Source	Estimate	Sq.root
S(Site)	50.886	7.1335
S(Year)	414.5	20.359
S(SitexYear)	-36.934	-6.0773
V(Res)	1162	34.088

Table N-11 PERMDISP test of 'Site' for edge habitat of upstream-downstream sites of Nepean River at the confluence of Matahil Creek into which West Camden WWTP discharges

DEVIATIONS FROM CENTROID

F: 0.2946 df1: 1 df2: 82

P(perm): 0.6072

MEANS AND STANDARD ERRORS

Group Size Average SE Downstream 49 38.341 0.68546 Upstream 35 38.913 0.79396



Table N-12 ANOSIM test of 'Site' period for edge habitat of upstream-downstream sites of Nepean River at the confluence of Matahil Creek into which West Camden WWTP discharges

Tests for differences between unordered Site period groups

Global Test

Sample statistic (R): 0.083

Significance level of sample statistic: 1.6%

Number of permutations: 9999 (Random sample from a large number)

Number of permuted statistics greater than or equal to R: 154

Pairwise Tests

	R	Significance	Possible	Actual	Number >=
Groups	Statistic	Level %	Permutations	Permutations	Observed
Downstream 1995 to 2018, Downstream 2018 to 2020	-0.098	75	211876	9999	7501
Downstream 1995 to 2018, Upstream 1995 to 2018	0.129	0.03	Very large	9999	2
Downstream 1995 to 2018, Upstream 2018 to 2020	-0.064	65.9	211876	9999	6591
Downstream 2018 to 2020, Upstream 1995 to 2018	0.068	30.6	52360	9999	3061
Downstream 2018 to 2020, Upstream 2018 to 2020	-0.063	60	35	35	21
Upstream 1995 to 2018, Upstream 2018 to 2020	0.081	27.9	52360	9999	2787





Subset of 66 (correlation 0.951) genera from edge habitat whose pattern matches that of the full set of 178 genera identified with the same subset found on two runs from 50 random start runs. Each run was based on three randomly selected genera. Genera were:

Aturidae Wheenyella, Coenagrionidae Ischnura, Physidae Physella, Chironomidae Cricotopus, Chironomidae Cryptochironomus, Chironomidae Dicrotendipes, Corbiculidae Corbicula, Dugesiidae Cura, Glossiphoniidae Helobdella, Lymnaeidae Pseudosuccinea, Planorbidae Gyraulus, Platycnemididae Nososticta, Pleidae Paraplea, Chironomidae Cladotanytarsus, Chironomidae Microtendipes, Chironomidae Parachironomus, Chironomidae Polypedilum, Chironomidae Procladius, Coenagrionidae Pseudagrion, Corduliidae Hemicordulia, Hydrophilidae Helochares, Isostictidae Rhadinosticta, Libellulidae Diplacodes, Libellulidae Nannophlebia, Limnesiidae Physolimnesia, Atyidae Paratya, Baetidae Cloeon, Caenidae Tasmanocoenis, Ceratopogonidae Bezzia, Ceratopogonidae Monohelea, Dytiscidae Necterosoma, Ecnomidae Ecnomus, Elmidae Coxelmis, Elmidae Ovolara, Gomphidae Austrogomphus, Leptoceridae Notalina, Corixidae Micronecta, Unionicolidae Unionicola, Chironomidae Coelopynia, Chironomidae Corynoneura, Chironomidae Parakiefferiella, Chironomidae Paramerina, Chironomidae Paratanytarsus, Chironomidae Riethia, Elmidae Hydora, Hydrodromidae Hydrodroma, Hydrophilidae Berosus, Leptoceridae Oecetis, Leptoceridae Triaenodes, Limnesiidae Limnesia, Mideopsidae Gretacarus, Oxidae Oxus, Unionicolidae Koenikea, Unionicolidae Recifella, Veliidae Microvelia, Calamoceratidae Anisocentropus, Chironomidae Ablabesmyia, Chironomidae Larsia, Chironomidae Tanytarsus, Dytiscidae Sternopriscus, Elmidae Austrolimnius, Hygrobatidae Coaustraliobates, Leptoceridae Triplectides, Oxidae Frontipoda, Ecnomidae Ecnomina, Stratiomyidae Odontomyia.






Figure N-9 Shade plot of freshwater macroinvertebrate edge habitat community structure of upstream-downstream sites of Nepean River at the confluence of Matahil Creek into which West Camden WWTP discharges





Stream health near Wallacia WWTP

The SIGNAL-SG plot for the Warragamba River provided an assessment of stream health. This plot was based on macroinvertebrate identification and counting results expressed as SIGNAL-SG scores and allows a visual comparison of data collected from 2019–20 against that collected between 2008 and 2019. This comparison suggests downstream stream health was maintained at a level comparable to that of the upstream site indicating wastewater discharge from Wallacia WWTP did not have a measurable impact on stream health during 2019-20 (Figure N-10).

A comparison of the upstream-downstream SIGNAL-SG scores for 2019-20 samples under a t-test returned a non-significant test outcome (Table N-14) and confirmed the visual trend.

As no measurable impact on downstream stream health was detected, no further data analysis was undertaken.




Figure N-10 Stream health of Warragamba River near Wallacia WWTP

Table N-14 t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from the Warragamba River near Wallacia WWTP

Method	Variances	DF	t Value	Pr > t
Pooled	Equal	12	0.76	0.4598

Equality of Variances					
Method	Num DF	Den DF	F Value	Pr > F	
Folded F	6	6	1.36	0.7176	

Stream health near Penrith WWTP

The SIGNAL-SG plots provided assessments of stream health for both Boundary Creek near Penrith WWTP and the Nepean River upstream-downstream of the confluence with Boundary Creek. These plots were based on macroinvertebrate identification and counting results expressed as SIGNAL-SG scores and allows a visual comparison of data collected from 2019–20 against that collected between 2003 to 2019 for the Boundary Creek sites and 1995 to 2019 for the Nepean River sites. These visual comparisons suggest downstream stream health was maintained at least at a level comparable to that of the upstream site indicating that the wastewater discharge from the Penrith WWTP did not have a measurable negative impact on stream health of either Boundary Creek (Figure N-11) or the Nepean River during 2019-20 (Figure N-12).

A comparison of the upstream-downstream SIGNAL-SG scores for 2019-20 samples under t-tests returned a non-significant test outcome for the Nepean River comparison (Table N-16) as well as the Boundary Creek comparison (Table N-15). As no measurable negative impact on downstream stream health was detected on either Boundary Creek or the Nepean River, no further data analysis was undertaken.




Figure N-11 Stream health of Boundary Creek near Penrith WWTP



Table N-15 t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from Boundary Creek near Penrith WWTP

Method	Variances	DF	t Value	Pr > t	
Pooled	Equal	10	1.70	0.1200	

Equality of Variances						
Method	Num DF	Den DF	F Value	Pr > F		
Folded F	5	5	1.77	0.5457		




Figure N-12 Stream health of the Nepean River upstream-downstream of the confluence of Boundary Creek near Penrith WWTP

Table N-16 t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from the Nepean River near Penrith WWTP

Method	Variances	DF	t Value	Pr > t
Pooled	Equal	10	-0.21	0.8408

Equality of Variances					
Method	Num DF	Den DF	F Value	Pr > F	
Folded F	5	5	1.11	0.9116	





The SIGNAL-SG plots provided assessments of stream health for both the unnamed creek near Winmalee WWTP and in the Nepean River situated upstream-downstream of the confluence with the unnamed creek. These plots were based on macroinvertebrate identification and counting results expressed as SIGNAL-SG scores and allows a visual comparison of data collected from 2019–20 against that collected between 2004 to 2019 for the unnamed creek and 1995 to 2019 for the Nepean River. These visual comparisons suggest downstream stream health was maintained in the Nepean River (Figure N-14) while stream health for the unnamed creek 0.3 km site remained at a relatively high level that overlapped the range of stream health observed for the 3 km downstream site that fell within the range observed over the 2004 to 2019 period (Figure N-13).

A comparison of the upstream-downstream SIGNAL-SG scores for 2019-20 samples under t-tests returned non-significant test outcomes for the Nepean River comparison (Table N-18) and the unnamed creek comparison (Table N-17).

As the non-significant t-test outcome for the unnamed creek was somewhat atypical to past year results, and an impact on unnamed creek has been routinely recorded in the past, further data analysis was undertaken.




Figure N-13 Stream health of unnamed creek below Winmalee WWTP for two downstream sites





Table N-17 t-test of both downstream site SIGNAL-SG scores from 2019-20 for unnamed creek below Winmalee WWTP

Method	Variances	DF	t Value	Pr > t
Pooled	Equal	10	1.52	0.1594

Equality of Variances					
Method	Num DF	Den DF	F Value	Pr > F	
Folded F	5	5	5.55	0.0833	




Figure N-14 Stream health of the Nepean River near Winmalee WWTP

Table N-18 t-test of upstream-downstream SIGNAL-SG scores from 2019-20 for the Nepean River near Winmalee WWTP

Method	Variances	DF	t Value	Pr > t
Pooled	Equal	14	-0.42	0.6773

Equality of Variances					
Method	Num DF	Den DF	F Value	Pr > F	
Folded F	7	7	1.17	0.8372	

As the unnamed creek has no flow upstream of Winmalee WWTP under dry weather conditions, both sampling sites were situated downstream of the WWTP. The first site is located 0.3 km downstream of the WWTP, while the second downstream site is situated 3 km downstream of the WWTP in a natural bushland catchment that lacks other anthropogenic influences. Both edge and riffle habitat data were collected consistently at both downstream sites on the same sampling occasions to allow multivariate analysis for the monitoring period of 2004 to 2020. Samples from each habitat were analysed separately.

Distinct groups of samples were evident in the three-dimensional ordination plot of edge habitat of the unnamed creek (Figure N-15). The n-MDS ordination pattern was confirmed in the corresponding tree diagram (Figure N-17) from classification analysis where the first and second divisions separated the 2018 to 2020 period samples, whereas the fourth division separated most 0.3 km downstream samples from most 3 km downstream samples (Figure N-17). Despite not showing the early separation between time periods, the riffle habitat showed a similar split between sites at around the third separation level in the corresponding tree diagram (Figure N-18) and clear grouping of sites in the three-dimensional ordination plot (Figure N-16).

The corresponding shade plots (Figure N-19 and Figure N-20) both displayed the tolerant taxon, the Blackfly larvae *Simulium* (SIGNAL-SG grade 4) as persistent through time and consistently abundant at the site 0.3 km downstream of the WWTP in both habitats. This taxon was absent on most collection occasions or occurred in much lower numbers at the 3 km downstream site. These shade plots illustrated that higher graded SIGNAL-SG taxa such as the non-biting midge larvae Chironomidae *Parametriocnemis* and caddisfly Leptoceridae *Triplectides* were more consistently collected from the site 3 km downstream, suggesting recovery in water quality with distance from the WWTP.

The BVSTEP routine was used to find a subset of taxa whose multivariate pattern matched that of the full dataset with 29 taxa (out of 91) identified for the edge habitat (Table N-27) and 20 taxa (out of 58) for the riffle habitat (Table N-28). These subsets of taxa form the main visual patterns in the respective shade plots (Figure N-19 and Figure N-20).

The PERMDISP analysis indicated a non-significant pattern of dispersion (spacing between same site samples) for the edge (Table N-23) and riffle (Table N-24) habitats. These results imply results of ANOSIM tests are focused on community structure differences between sites.

An ANOSIM test was run on the factor 'Site'. The returned ANOSIM R-values were at mid-range levels (Table N-19) and (Table N-20), implying both downstream sites assemblage structures were distinguishable for both habitats.

To further explore the community structure, hypothesis testing was conducted with a PERMANOVA model. This model comprised the fixed factors 'Site' and 'Year'. 'Year' represented samples collected in years between 2004 and 2020 and 'Site' having two levels, 0.3 km downstream and 3 km downstream. A statistically non-significant 'Site x Year' interaction was returned for the edge (Table N-21) and riffle (Table N-22) habitats. These non-significant results allowed us to view the 'Site' and 'Year' results. Statistically significant results were returned for 'Year' and 'Site' factors. The estimates of components of variation indicated 'Site' explained approximately twice the variation than that explained by 'Year' for the edge habitat (Table N-21) and four times the variation than that explained by 'Year' for the riffle habitat (Table N-22).

A second run of ANOSIM based on 'Site-period' groups displayed in ordination plots Figure N-15 and Figure N-16 returned a significant global mid-range R-value of 0.60 for the edge habitat. In the resulting pairwise comparisons, four tests returned R-values at a level (R = 1.0, 0.875, 0.838 and

0.782, Table N-25) that can be expected from natural differences between groups from variation in the substratum composition of the habitats between sites. Besley and Chessman (2008) found R-values up to 0.66 for sites on the same near-pristine stream. A lower mid-range global R-value of 0.429 was returned for the riffle habitat with only a single corresponding pairwise test for the riffle habitat returned above an R-value of 0.66 (Table N-26).

These multivariate analysis results suggested community structure alteration from wastewater discharge in the unnamed creek was most evident in macroinvertebrate assemblages within the edge habitat.








Figure N-15 Dimensions 1 and 3 of three-dimensional ordination plot of freshwater macroinvertebrate edge habitat community structure of both downstream sites of unnamed creek below Winmalee WWTP




Figure N-16 Dimensions 1 and 2 of three-dimensional ordination plot of freshwater macroinvertebrate riffle habitat community structure of both downstream sites of unnamed creek below Winmalee WWTP








Figure N-17 Tree diagram of freshwater macroinvertebrate edge habitat community structure of both downstream sites of unnamed creek below Winmalee WWTP




Figure N-18 Tree diagram of freshwater macroinvertebrate riffle habitat community structure of both downstream sites of unnamed creek below Winmalee WWTP





Table N-19 ANOSIM test of 'Site' factor for edge habitat unnamed creek near Winmalee WWTP

Tests for differences between unordered Site groups

Global Test

Sample statistic (R): 0.538

Significance level of sample statistic: 0.01%

Number of permutations: 9999 (Random sample from a large number)

Number of permuted statistics greater than or equal to R: 0

Table N-20 ANOSIM test of 'Site' factor for riffle habitat unnamed creek below Winmalee WWTP

Tests for differences between unordered Site groups

Global Test

Sample statistic (R): 0.49

Significance level of sample statistic: 0.01%

Number of permutations: 9999 (Random sample from a large number)

Number of permuted statistics greater than or equal to R: 0





Table N-21 PERMANOVA test of 'Site' and 'Year' factors for edge habitat unnamed creek below Winmalee WWTP

Sums of squares type: Type III (partial) Fixed effects sum to zero for mixed terms

Permutation method: Permutation of residuals under a reduced model

Number of permutations: 9999

PERMANOVA table of results

						Unique
Source	df	SS	MS	Pseudo-F	P(perm)	perms
Site	1	20821	20821	14.671	0.0001	9937
Year	16	43702	2731.4	1.9246	0.0001	9749
SitexYear	16	26584	1661.5	1.1707	0.0601	9742
Res	30	42576	1419.2			
Total 63	1.3	3517E+05				

Estimates of components of variation

Estimate	Sq.root
637.77	25.254
349.19	18.687
128.94	11.355
1419.2	37.672
	Estimate 637.77 349.19 128.94

Table N-22 PERMANOVA test of 'Site' and 'Year' factors for riffle habitat unnamed creek below Winmalee WWTP

Sums of squares type: Type III (partial)
Fixed effects sum to zero for mixed terms

Permutation method: Permutation of residuals under a reduced model

Number of permutations: 9999

PERMANOVA table of results

						Unique
Source	df	SS	MS	Pseudo-F	P(perm)	perms
Site	1	19857	19857	17.595	0.0001	9934
Year	16	26825	1676.6	1.4856	0.0031	9797
SitexYear	16	16217	1013.6	0.89811	0.7454	9811
Res	30	33857	1128.6			
Total	63	96518				

Estimates of components of variation

Source	Estimate	Sq.root
S(Site)	615.63	24.812
S(Year)	145.83	12.076
S(SitexYear)	-61.201	-7.8231
V(Res)	1128.6	33.594







DEVIATIONS FROM CENTROID

F: 0.29336 df1: 1 df2: 62

P(perm): 0.6204

MEANS AND STANDARD ERRORS

 Group
 Size
 Average
 SE

 3km downstream
 32
 41.868
 1.1904

 0.3km downstream
 32
 40.881
 1.3777

Table N-24 PERMDISP test of 'Site' factor for riffle habitat unnamed creek below Winmalee WWTP

DEVIATIONS FROM CENTROID

F: 0.45791 df1: 1 df2: 62

P(perm): 0.5305

MEANS AND STANDARD ERRORS

Group	Size	Average	SE
3km downstream	32	34.468	1.2517
0.3km downstream	32	33.163	1.4666





Table N-25 ANOSIM test of 'Site period' samples for edge habitat unnamed creek below Winmalee WWTP

Global Test

Sample statistic (R): 0.6

Significance level of sample statistic: 0.01%

Number of permutations: 9999 (Random sample from a large number)

Number of permuted statistics greater than or equal to R: 0

Pairwise Tests

	R	Significance	Possible	Actual	Number >=
Groups	Statistic	Level %	Permutations	Permutations	Observed
3km downstream 2004 to 2018, 3km downstream 2018 to 2020	0.116	21.3	35960	9999	2133
3km downstream 2004 to 2018, 0.3km downstream 2004 to 2018	0.562	0.01	Very large	9999	0
3km downstream 2004 to 2018, 0.3km downstream 2018 to 2020	0.875	0.01	35960	9999	0
3km downstream 2018 to 2020, 0.3km downstream 2004 to 2018	0.838	0.02	35960	9999	1
3km downstream 2018 to 2020, 0.3km downstream 2018 to 2020	1	2.9	35	35	1
0.3km downstream 2004 to 2018, 0.3km downstream 2018 to 2020	0.782	0.02	35960	9999	1

Table N-26 ANOSIM test of 'Site period' samples for riffle habitat unnamed creek below Winmalee WWTP

Global Test

Sample statistic (R): 0.429

Significance level of sample statistic: 0.01%

Number of permutations: 9999 (Random sample from a large number)

Number of permuted statistics greater than or equal to R: 0

Pairwise Tests

	R	Significance	Possible	Actual	Number >=
Groups	Statistic	Level %	Permutations	Permutations	Observed
Downstream 2004 to 2018, Downstream 2018 to 2020	0.225	7.8	35960	9999	776
Downstream 2004 to 2018, 0.3km downstream 2004 to 2018	0.493	0.01	Very large	9999	0
Downstream 2004 to 2018, 0.3km downstream 2018 to 2020	0.441	0.5	35960	9999	51
Downstream 2018 to 2020, 0.3km downstream 2004 to 2018	0.635	0.05	35960	9999	4
Downstream 2018 to 2020, 0.3km downstream 2018 to 2020	0.771	2.9	35	35	1
0.3km downstream 2004 to 2018, 0.3km downstream 2018 to 2020	-0.083	66.1	35960	9999	6606



Table N-27 Genera subset whose multivariate pattern matches full genera set of the edge habitat unnamed creek below Winmalee WWTP Subset of 29 (correlation 0.951) genera from edge habitat whose pattern matches that of the full set of 91 genera identified with the same subset found on 22 runs from 50 random start runs. Each run was based on three randomly selected genera. Genera were:

Chironomidae *Chironomus*, Physidae *Physella*, Chironomidae *Cricotopus*, Dugesiidae *Cura*, Glossiphoniidae *Helobdella*, Lumbriculidae *Lumbriculus*, Simuliidae *Simulium*, Argiolestidae *Austroargiolestes*, Chironomidae *Microtendipes*, Chironomidae *Polypedilum*, Chironomidae *Rheocricotopus*, Chironomidae *Rheotanytarsus*, Chironomidae *Thienemanniella*, Hydropsychidae *Cheumatopsyche*, Libellulidae *Nannophlebia*, Scyphacidae *Haloniscus*, Talitridae *Arcitalitrus*, Tateidae *Potamopyrgus*, Atyidae *Paratya*, Elmidae *Notriolus*, Elmidae *Simsonia*, Hydraenidae *Hydraena*, Hydroptilidae *Hellyethira*, Leptoceridae *Notalina*, Micronectidae *Micronecta*, Aeshnidae *Austroaeschna*, Veliidae *Microvelia*, Chironomidae *Tanytarsus*, Leptoceridae *Triplectides*.

Table N-28 Genera subset whose multivariate pattern matches full genera set of the riffle habitat unnamed creek below Winmalee WWTP Subset of 20 (correlation 0.954) genera from riffle habitat whose pattern matches that of the full set of 58 genera identified with the same subset found on one run from 50 random start runs. Each run was based on three randomly selected genera. Genera were:

Chironomidae *Chironomus*, Chironomidae *Cardiocladius*, Chironomidae *Cricotopus*, Dugesiidae *Cura*, Naididae *Nais*, Simuliidae *Simulium*, Argiolestidae *Austroargiolestes*, Chironomidae *Eukiefferiella*, Chironomidae *Polypedilum*, Chironomidae *Rheocricotopus*, Chironomidae *Rheotanytarsus*, Chironomidae *Thienemanniella*, Hydropsychidae *Cheumatopsyche*, Libellulidae *Nannophlebia*, Tateidae *Potamopyrgus*, Elmidae *Notriolus*, Elmidae *Simsonia*, Hydrobiosidae *Ulmerochorema*, Aeshnidae *Austroaeschna*, Chironomidae *Parametriocnemus*.








Figure N-19 Shade plot of freshwater macroinvertebrate edge habitat community structure of both downstream sites of unnamed creek below Winmalee WWTP




Figure N-20 Shade plot of freshwater macroinvertebrate riffle habitat community structure of both downstream sites of unnamed creek below Winmalee WWTP

Sufficient macrophyte and edge habitat data were collected consistently enough at upstream-downstream sites on the same sampling occasions to allow multivariate analysis for the monitoring period of 1995 to 2020 (less sample collection gaps outlined in Volume 1 Table 3-2). Samples from each habitat were analysed separately.

The Nepean River macrophyte and edge habitat data pattern were visually displayed in a three-dimensional nMDS ordination plots achieve an acceptable level of fit (stress) due to inherent variation. Data points were colour coded by 'Site period' with two periods 1995 to 2018 and 2018 to 2020. There was no clear separation of groups of upstream and downstream samples in either of the ordination plots (Figure N-21 and Figure N-22). Rather a mix of upstream and downstream samples was observed, with most recent samples intermingling with past samples.

The lack of a clear pattern between sites in the ordination plots (Figure N-21 and Figure N-22) was also apparent in the corresponding tree diagrams (Figure N-23 and Figure N-24) and shade plots (Figure N-25 and Figure N-26) suggesting communities between sites were similar. Subsets of taxa defining the multivariate pattern are listed in Table N-37 and Table N-38.

The PERMDISP analysis returned non-significant for both macrophyte (Table N-33) and edge (Table N-34) habitats. This implies results of ANOSIM tests are focused on community structure differences between upstream-downstream sites.

An ANOSIM test was run on the factor 'Site'. The returned ANOSIM R-values were at very low levels close to zero (Table N-29 and Table N-30) implying the assemblage structure of sites were almost indistinguishable.

To further explore community structure, hypothesis testing was conducted with a PERMANOVA model. This model comprised the fixed factors 'Site' and 'Year' with 'Year' representing samples collected in years between 1995 and 2020 and 'Site' having two levels, upstream and downstream. For the macrophyte habitat a statistically non-significant 'Site x Year' interaction was returned (Table N-31). This non-significant result allowed us to view the 'Site' and 'Year' results. Significant results were returned for 'Site' and 'Year' factors. Inspecting estimates of components of variation output indicated that 'Year' explained about seven times the variation than that explained by 'Site' (Table N-31). For the edge habitat, a statistically non-significant 'Site x Year' interaction was returned (Table N-32). Both 'Site' and 'Year' factors resulted in significant results. Inspecting estimates of components of variation output indicated 'Year' explained more than three times the variation than that explained by 'Site' (Table N-32).

A second run of ANOSIM based on 'Site-period' samples displayed in ordination plots (Figure N-21 and Figure N-22) returned a non-significant global R-value for the macrophyte habitat (Table N-35) and for the edge habitat a significant global R-value of 0.104, but this was relatively low (Table N-36). Inspection of pairwise tests for the edge habitat indicated five of the six comparisons were non-significant and the only significant test had a low-range R-value of 0.111 (Table N-36).

These results suggested community structure in the unnamed creek near the WWTP was altered by wastewater discharge from Winmalee WWTP but this impact did not extend as far as the Nepean River.








Figure N-21 Dimensions 1 and 3 of three-dimensional nMDS ordination plot of freshwater macroinvertebrate macrophyte habitat community structure of sites upstream-downstream of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges




Figure N-22 Dimensions 1 and 3 of three-dimensional ordination plot of freshwater macroinvertebrate edge habitat community structure of sites upstream-downstream of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges








Figure N-23 Tree diagram of freshwater macroinvertebrate macrophyte habitat community structure of sites upstream-downstream of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges




Figure N-24 Tree diagram of freshwater macroinvertebrate edge habitat community structure of sites upstream-downstream of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges





Table N-29 ANOSIM test of 'Site' for macrophyte habitat of upstream-downstream sites of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges

Tests for differences between unordered Site groups

Global Test

Sample statistic (R): 0.035

Significance level of sample statistic: 2.1%

Number of permutations: 9999 (Random sample from a large number)

Number of permuted statistics greater than or equal to R: 210

Table N-30 ANOSIM test of 'Site' for edge habitat Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges

Tests for differences between unordered Site groups

Global Test

Sample statistic (R): 0.101

Significance level of sample statistic: 0.01%

Number of permutations: 9999 (Random sample from a large number)

Number of permuted statistics greater than or equal to R: 0

Table N-31 PERMANOVA test of 'Site' and 'Year' factors for macrophyte habitat of upstream-downstream sites of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges

Sums of squares type: Type III (partial) Fixed effects sum to zero for mixed terms

Permutation method: Permutation of residuals under a reduced model

Number of permutations: 9999 *PERMANOVA table of results*

		41010 01 1000				Unique			
Source	df	SS	MS	Pseudo-F	P(perm)	perms			
Site	1	3834.8	3834.8	2.2594	0.0074	9916			
Year	25	74831	2993.3	1.7636	0.0001	9711			
SitexYear	25	40407	1616.3	0.95227	0.7067	9700			
Res	45	76378	1697.3						
Total	96	1.9655E+0)5						
Estimates of components of variation									
Source		Estimate	Sq.root						
S(Site)		46.64	6.8294						
S(Year)		351.42	18.746						
S(SitexYear	.)	-43.939	-6.6286						
V(Res)		1697.3	41.198						

Table N-32 PERMANOVA test of 'Site' and 'Year' factors for edge habitat of upstreamdownstream sites of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges

Sums of squares type: Type III (partial) Fixed effects sum to zero for mixed terms

Permutation method: Permutation of residuals under a reduced model

Number of permutations: 9999 PERMANOVA table of results

							Unique
Source	df	SS		MS	Pseudo-F	P(perm)	perms
Site	1	5642.	2	5642.2	3.3819	0.0001	9920
Year	25	7129	6	2851.9	1.7094	0.0001	9638
SitexYear**	24	4302	4	1792.6	1.0745	0.1634	9638
Res	45	7507	7	1668.4			
Total	95	1.9566E	+05				
Estimates of components of variation							
Source	Es	timate	Sq.	root			
S(Site)	8	7.425	9.3	501			
S(Year)	3	24.51	18.	014			
S(SitexYear)	6	6.165	8.1	342			
V(Res)	1	668.4	40.	846			




Table N-33 PERMDISP test of 'Site' for macrophyte habitat of upstreamdownstream sites of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges

DEVIATIONS FROM CENTROID

F: 0.098726 df1: 1 df2: 95

P(perm): 0.762

MEANS AND STANDARD ERRORS

 Group
 Size
 Average
 SE

 Downstream
 47
 43.744
 0.97847

 Upstream
 50
 44.192
 1.0318

Table N-34 PERMDISP test of 'Site' for edge habitat of upstream-downstream sites of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges

DEVIATIONS FROM CENTROID

F: 0.015608 df1: 1 df2: 94

P(perm): 0.9019

MEANS AND STANDARD ERRORS

 Group
 Size
 Average
 SE

 Downstream
 48
 44.123
 0.83037

 Upstream
 48
 43.982
 0.76377



Table N-35 ANOSIM test of 'Site period' for macrophyte habitat of upstream-downstream sites of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges

Tests for differences between unordered Site period groups

Global Test

Sample statistic (R): 0.025

Significance level of sample statistic: 19.8%

Number of permutations: 9999 (Random sample from a large number)

Number of permuted statistics greater than or equal to R: 1977

Pairwise Tests

R	Significance	Possible	Actual	Number >=
Statistic	Level %	Permutations	Permutations	Observed
-0.039	60	178365	9999	5997
0.037	2.3	Very large	9999	229
0.105	19.9	178365	9999	1985
-0.107	76.9	230300	9999	7687
0.302	8.6	35	35	3
-0.006	48.2	230300	9999	4822
	Statistic -0.039 0.037 0.105 -0.107 0.302	Statistic Level % -0.039 60 0.037 2.3 0.105 19.9 -0.107 76.9 0.302 8.6	Statistic Level % Permutations -0.039 60 178365 0.037 2.3 Very large 0.105 19.9 178365 -0.107 76.9 230300 0.302 8.6 35	Statistic Level % Permutations Permutations -0.039 60 178365 9999 0.037 2.3 Very large 9999 0.105 19.9 178365 9999 -0.107 76.9 230300 9999 0.302 8.6 35 35

Table N-36 ANOSIM test of 'Site period' for edge habitat of upstream-downstream sites of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges

Tests for differences between unordered Site period groups

Global Test

Sample statistic (R): 0.104

Significance level of sample statistic: 0.04%

Number of permutations: 9999 (Random sample from a large number)

Number of permuted statistics greater than or equal to R: 3

Pairwise Tests

	R	Significance	Possible	Actual	Number >=
Groups	Statistic	Level %	Permutations	Permutations	Observed
Downstream 1995 to 2018, Downstream 2018 to 2020	-0.073	68.8	194580	9999	6875
Downstream 1995 to 2018, Upstream 1995 to 2018	0.111	0.01	Very large	9999	0
Downstream 1995 to 2018, Upstream 2018 to 2020	0.206	8.3	194580	9999	832
Downstream 2018 to 2020, Upstream 1995 to 2018	0.014	42.7	194580	9999	4264
Downstream 2018 to 2020, Upstream 2018 to 2020	0.26	17.1	35	35	6
Upstream 1995 to 2018, Upstream 2018 to 2020	0.183	9.8	194580	9999	979





Table N-37 Genera subset whose multivariate pattern matches full genera set of the macrophyte habitat of upstream-downstream sites of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges

Subset of 34 (correlation 0.950) genera from macrophyte habitat whose pattern matches that of the full set of 129 genera identified with the same subset found on two runs from 50 random start runs. Each run was based on three randomly selected genera. Genera were:

Coenagrionidae *Ischnura*, Physidae *Physella*, Chironomidae *Cricotopus*, Chironomidae *Dicrotendipes*, Corbiculidae *Corbicula*, Dugesiidae *Cura*, Lymnaeidae *Pseudosuccinea*, Planorbidae *Ferrissia*, Simuliidae *Simulium*, Chironomidae *Cladotanytarsus*, Chironomidae *Microtendipes*, Chironomidae *Polypedilum*, Chironomidae *Rheotanytarsus*, Chironomidae *Thienemanniella*, Coenagrionidae *Pseudagrion*, Corduliidae *Hemicordulia*, Hebridae *Merragata*, Hydrophilidae *Helochares*, Hydropsychidae *Cheumatopsyche*, Libellulidae *Diplacodes*, Libellulidae *Nannophlebia*, Limnesiidae *Physolimnesia*, Atyidae *Paratya*, Baetidae *Cloeon*, Caenidae *Tasmanocoenis*, Ceratopogonidae *Bezzia*, Ecnomidae *Ecnomus*, Hydroptilidae *Hellyethira*, Leptoceridae *Notalina*, Baetidae *Offadens*, Hydrodromidae *Hydrodroma*, Leptoceridae *Oecetis*, Veliidae *Microvelia*, Chironomidae *Tanytarsus*.

Table N-38 Genera subset whose multivariate pattern matches full genera set of the edge habitat of upstream-downstream sites of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges

Subset of 51 (correlation 0.951) genera from edge habitat whose pattern matches that of the full set of 146 genera identified with the same subset found on one run from 50 random start runs. Each run was based on three randomly selected genera. Genera were:

Chironomidae Chironomus, Coenagrionidae Ischnura, Physidae Physella, Chironomidae Cricotopus, Chironomidae Cryptochironomus, Chironomidae Dicrotendipes, Coenagrionidae Austroagrion, Corbiculidae Corbicula, Dugesiidae Cura, Glossiphoniidae Helobdella, Lumbriculidae Lumbriculus, Lymnaeidae Pseudosuccinea, Naucoridae Naucoris, Planorbidae Gyraulus, Platycnemididae Nososticta, Simuliidae Simulium, Chironomidae Cladotanytarsus, Chironomidae Microtendipes, Chironomidae Polypedilum, Chironomidae Procladius, Chironomidae Rheocricotopus, Chironomidae Rheotanytarsus, Coenagrionidae Pseudagrion, Hydrophilidae Helochares, Hydropsychidae Cheumatopsyche, Libellulidae Diplacodes, Libellulidae Nannophlebia, Limnesiidae Physolimnesia, Mesoveliidae Mesovelia, Atyidae Paratya, Baetidae Cloeon, Caenidae Tasmanocoenis, Ceratopogonidae Bezzia, Ecnomidae Ecnomus, Gomphidae Austrogomphus, Hydroptilidae Hellyethira, Leptoceridae Notalina, Libellulidae Orthetrum, Micronectidae Micronecta, Baetidae Offadens, Chironomidae Corynoneura, Hydrodromidae Hydrodroma, Hydrophilidae Berosus, Leptoceridae Oecetis, Unionicolidae Recifella, Veliidae Microvelia, Calamoceratidae Anisocentropus, Chironomidae Ablabesmyia, Chironomidae Tanytarsus, Hygrobatidae Coaustraliobates, Leptoceridae Triplectides.








Figure N-25 Shade plot of freshwater macroinvertebrate macrophyte habitat community structure of upstream-downstream sites of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges






Figure N-26 Shade plot of freshwater macroinvertebrate edge habitat community structure of upstream-downstream sites of Nepean River at the confluence of the unnamed creek into which Winmalee WWTP discharges





Stream health near North Richmond WWTP

The SIGNAL-SG plots provided assessments of stream health for both Redbank Creek near North Richmond WWTP and in the Hawkesbury River upstream-downstream of the confluence with Redbank Creek. These plots were based on macroinvertebrate identification and counting results expressed as SIGNAL-SG scores and allows a visual comparison of data collected from 2019–20 against that collected between 2004 to 2019 for Redbank Creek and 1995 to 2019 for the Hawkesbury River. These visual comparisons suggest that the wastewater discharges from the North Richmond WWTP did not have a measurable negative impact on stream health of the Hawkesbury River over 1995 to 2020 (Figure N-27).

A comparison of the upstream-downstream SIGNAL-SG scores for 2019-20 samples under t-tests returned non-significant outcomes for both the Hawkesbury River and Redbank Creek (Table N-39 and Table N-40) and confirmed the visual trend for 2019-20.

As no measurable impact on downstream stream health was detected, no further data analysis was undertaken.




Figure N-27 Stream health of Hawkesbury River upstream-downstream of the confluence of Redbank Creek near North Richmond WWTP





Table N-39 t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from the Hawkesbury River near North Richmond WWTP

Method	Variances	DF	t Value	Pr > t
Pooled	Equal	10	1.77	0.1074

Equality of Variances					
Method	Num DF	Den DF	F Value	Pr > F	
Folded F	5	5	3.12	0.2371	




Figure N-28 Stream health of Redbank Creek near North Richmond WWTP

Table N-40 t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from Redbank Creek near North Richmond WWTP

Method	Variances	DF	t Value	Pr > t
Pooled	Equal	10	1.00	0.3422

Equality of Variances					
Method Num DF Den DF F Value Pr > F					
Folded F	5	5	1.77	0.5465	





Stream health near St Mary's WWTP

The SIGNAL-SG plot for South Creek provided an assessment of stream health. This plot was based on macroinvertebrate identification and counting results expressed as SIGNAL-SG scores and allows a visual comparison of data collected from 2019–20 against that collected between 1995 and 2019. This comparison suggests downstream stream health was maintained at a level comparable to that of the upstream site indicating wastewater discharge from St Mary's WWTP did not have a measurable impact on stream health during 2019-20 (Figure N-29). When viewing historical aerial maps on Nearmap, it is noted there was a reduction in land clearing activities immediately north of the upstream site in 2017-18 to 2019-20. This may be contributing to the overall improved stream health.

A comparison of the upstream-downstream SIGNAL-SG scores for 2019-20 samples under a t-test returned a non-significant test outcome (Table N-41) and confirmed the visual trend of the SIGNAL-SG plot.

As no measurable negative impact on downstream stream health was detected, no further data analysis was undertaken.




Figure N-29 Stream health of South Creek near St Mary's WWTP

Table N-41 t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from South Creek near St Mary's WWTP

Method	Variances	DF	t Value	Pr > t	
Pooled	Equal	10	-2.03	0.0695	

Equality of Variances					
Method Num DF Den DF F Value Pr > F					
Folded F	5	5	2.11	0.4317	





Stream health near Quakers Hill WWTP

The SIGNAL-SG plot for Breakfast Creek provided an assessment of stream health. This plot was based on macroinvertebrate identification and counting results expressed as SIGNAL-SG scores and allows a visual comparison of data collected from 2019–20 against that collected between 1995 and 2019. This comparison suggests downstream stream health was maintained at a level comparable to that of the upstream site indicating wastewater discharge from Quakers Hill WWTP did not have a measurable impact on stream health during 2019-20 (Figure N-30).

A comparison of the upstream-downstream SIGNAL-SG scores for 2019-20 samples under a t-test returned a non-significant test outcome (Table N-42) and confirmed the visual trend of the SIGNAL-SG plot.

As no measurable impact on downstream stream health was detected, no further data analysis was undertaken.




Figure N-30 Stream health of Breakfast Creek near Quakers Hill WWTP

Table N-42 t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from Breakfast Creek near Quakers Hill WWTP

Method	Variances	DF	t Value	Pr > t
Pooled	Equal	10	1.33	0.2118

Equality of Variances					
Method Num DF Den DF F Value Pr > F					
Folded F	5	5	3.33	0.2125	





Stream health near Riverstone WWTP

The SIGNAL-SG plot for Eastern Creek provided an assessment of stream health. This plot was based on macroinvertebrate identification and counting results expressed as SIGNAL-SG scores and allows a visual comparison of data collected from 2019–20 against that collected between 1995 and 2019. This comparison suggests downstream stream health was maintained at a level comparable to that of the upstream site indicating wastewater discharge from Riverstone WWTP did not have a measurable impact on stream health during 2019-20 (Figure N-31).

A comparison of the upstream-downstream SIGNAL-SG scores for 2019-20 samples under a t-test returned a non-significant test outcome (Table N-43) and confirmed the visual trend.

As no measurable negative impact on downstream stream health was detected, no further data analysis was undertaken.




Figure N-31 Stream health of Eastern Creek near Riverstone WWTP

Table N-43 t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from Eastern Creek near Riverstone WWTP

Method	Variances	DF	t Value	Pr > t
Pooled	Equal	14	1.84	0.0878

Equality of Variances					
Method Num DF Den DF F Value Pr > F					
Folded F	7	7	3.20	0.1476	





Stream health near Castle Hill WWTP

The SIGNAL-SG plot for Cattai Creek provided an assessment of stream health. This plot was based on macroinvertebrate identification and counting results expressed as SIGNAL-SG scores and allows a visual comparison of data collected from 2019–20 against that collected between 1995 and 2019. This comparison suggests downstream stream health was maintained at a level comparable to that of the upstream site indicating wastewater discharge from Castle Hill WWTP did not have a measurable impact on stream health during 2019-20 (Figure N-32).

A comparison of the upstream-downstream SIGNAL-SG scores for 2019-20 samples under a t-test returned a non-significant test outcome (Table N-44) and confirmed the visual trend.

As no measurable impact on downstream stream health was detected, no further data analysis was undertaken.




Figure N-32 Stream health of Cattai Creek near Castle Hill WWTP

Table N-44 t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from Cattai Creek near Castle Hill WWTP

Method	Variances	DF	t Value	Pr > t
Pooled	Equal	10	-1.67	0.1267

Equality of Variances					
Method Num DF Den DF F Value Pr > F					
Folded F	5	5	2.01	0.4627	





Stream health near Rouse Hill WWTP

The SIGNAL-SG plot for Second Ponds Creek provided an assessment of stream health. This plot was based on macroinvertebrate identification and counting results expressed as SIGNAL-SG scores and allows a visual comparison of data collected from 2019–20 against that collected between 1995 and 2019. This comparison suggests downstream stream health was maintained at a level comparable to that of the upstream site indicating wastewater discharge from Rouse Hill WWTP did not have a measurable impact on stream health during 2019-20 (Figure N-33).

A comparison of the upstream-downstream SIGNAL-SG scores for 2019-20 samples under a t-test returned a non-significant test outcome (Table N-45) and confirmed the visual trend.

As no measurable negative impact on downstream stream health was detected, no further data analysis was undertaken.




Figure N-33 Stream health of Second Ponds Creek near Rouse Hill WWTP

Table N-45 t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from Second Ponds Creek near Rouse Hill WWTP

Method	Variances	DF	t Value	Pr > t
Satterthwaite	Unequal	6.292	-1.81	0.1174

Equality of Variances					
Method	Num DF	Den DF	F Value	Pr > F	
Folded F	5	5	7.61	0.0439	





Stream health near Hornsby Heights WWTP

The SIGNAL-SG plot for Calna Creek provided an assessment of stream health. This plot was based on macroinvertebrate identification and counting results expressed as SIGNAL-SG scores and allows a visual comparison of data collected from 2019–20 against that collected between 1996 and 2019. This comparison suggests downstream stream health has not been maintained at a level comparable to that of the upstream site, indicating wastewater discharge from the Hornsby Heights WWTP did have a measurable persistent impact on stream health during over the last nine financial years (Figure N-34).

A comparison of the upstream-downstream SIGNAL-SG scores for 2019-20 samples under a t-test returned a significant test outcome (Table N-46).

As a measurable negative impact on downstream stream health was detected, further analysis of the macroinvertebrate community data was undertaken to explore for trends not revealed in the SIGNAL-SG analysis.




Figure N-34 Stream health of Calna Creek near Hornsby Heights WWTP

Table N-46 t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from Calna Creek near Hornsby Heights WWTP

Method	Variances	DF	t Value	Pr > t
Pooled	+Equal	10	-4.93	0.0006

Equality of Variances					
Method	Num DF	Den DF	F Value	Pr > F	
Folded F	5	5	2.50	0.3378	



Both edge and riffle habitats were collected consistently enough at upstream-downstream sites on the same sampling occasions to allow a multivariate analysis for the monitoring period of 1996 to 2020. Each habitat (edge and riffle) was analysed separately with comparisons assessed with upstream-downstream sites.

In the three-dimensional nMDS ordination plot of the Calna Creek edge habitat, a relatively interspersed pattern of upstream and downstream samples were observed (Figure N-35). This pattern was confirmed in the corresponding tree diagram from cluster analysis as the first division did not separate a group of upstream samples from another group of downstream samples (Figure N-37). The riffle habitat pattern displayed less overlap of upstream-downstream samples in the Calna Creek ordination plot (Figure N-36) and tree diagram (Figure N-38) compared to the edge habitat.

The PERMDISP analysis indicated a similar pattern of dispersion (spacing between same samples) for the upstream and downstream sites of the riffle habitat (Table N-52). This suggests the variability in taxonomic composition of samples collected over time was similar for upstream and downstream riffle sites through the period 1995 to 2020. As such, the subsequent riffle habitat results of ANOSIM tests were focused on community structure differences between sites. In contrast, significant dispersion was shown for the edge habitat samples (Table N-51). This outcome suggests subsequent edge habitat results of ANOSIM tests are describing both the variability in taxonomic composition of samples over time as well as community composition variability between the upstream and downstream sites.

An ANOSIM test was run on the factor 'Site'. The returned R-values were at a low-range level for edge (Table N-47) and at a mid-range level for riffle (Table N-48). These R-value results suggest site specific assemblages were more distinguishable for the riffle habitat and less distinguishable for the edge habitat. This pattern is reinforced by the shade plots that show a clear difference in sites within the riffle habitat (Figure N-40) and a less distinct pattern within the edge habitat (Figure N-39). These shade plots also show the riffle habitat has a smaller set of taxa (105) compared with the more diverse edge habitat (139 taxa). The BVSTEP routine was used to find a subset of taxa whose multivariate pattern matched that of the full dataset with 30 taxa identified for the riffle habitat (Table N-56) and 36 taxa for the edge habitat (Table N-55). These subsets of taxa reflect those taxa which formed the main visual patterns in the respective shade plots.

To further explore community structure, hypothesis testing was conducted with a PERMANOVA model. This model comprised the fixed factors 'Site' and 'Year' with 'Year' representing samples collected between 1996 and 2020 and 'Site' having two levels, upstream and downstream. A statistically significant 'Site x Year' interaction was returned for both the edge and the riffle habitats (Table N-49 and Table N-50) suggesting a change through time at least at one site.

A second run of ANOSIM based on 'Site-Period' groups in the 3D ordination plots (Figure N-35 and Figure N-36) returned a significant global low-range R-value (0.299) for the edge habitat. Under subsequent upstream-downstream pairwise comparisons, two tests returned an R-value at a level (R = 0.948 and 0.851) (Table N-53) that were not expected from natural differences between groups from variation in the substratum composition of the habitats between sites. Besley and Chessman (2008) found R-values up to 0.66 for sites on the same near-pristine stream. A mid-range global R-value (0.579) was returned for the riffle habitat and two of the upstream and downstream pairwise comparisons returned R-values (0.936 and 0.823) (Table N-54) that were at a level that implied more than natural substratum differences between sites.

In summary, the SIGNAL-SG control chart plot showed clear differences between the upstream-downstream sites consistently over the last nine financial years. Variability in the range of stream health levels were also evident for upstream-downstream sites in this SIGNAL-SG control chart. This variability and difference in assemblage structure suggested by SIGNAL-SG results was also evident in multivariate analysis. Both SIGNAL-SG and multivariate results suggest downstream community structure in Calna Creek has been consistently altered by wastewater discharge from the Hornsby Heights WWTP through the 2011 to 2020 monitoring period.




Figure N-35 Dimensions 1 and 3 of three-dimensional ordination plot of freshwater macroinvertebrate edge habitat community structure of upstream-downstream sites of Calna Creek near Hornsby Heights WWTP








Figure N-36 Dimensions 1 and 2 of three-dimensional ordination plot of freshwater macroinvertebrate riffle habitat community structure of upstream-downstream sites of Calna Creek near Hornsby Heights WWTP




Figure N-37 Tree diagram of freshwater macroinvertebrate edge habitat community structure of upstream-downstream sites of Calna Creek near Hornsby Heights WWTP








Figure N-38 Tree diagram of freshwater macroinvertebrate riffle habitat community structure of upstream-downstream sites of Calna Creek near Hornsby Heights WWTP

Table N-47 ANOSIM test of 'Site' factor for edge habitat of Calna Creek near Hornsby Heights www.TP

Tests for differences between unordered Site groups

Global Test

Sample statistic (R): 0.36

Significance level of sample statistic: 0.01%

Number of permutations: 9999 (Random sample from a large number)

Number of permuted statistics greater than or equal to R: 0

Table N-48 ANOSIM test of 'Site' factor for riffle habitat of Calna Creek near Hornsby Heights WWTP

Tests for differences between unordered Site groups

Global Test

Sample statistic (R): 0.586

Significance level of sample statistic: 0.01%

Number of permutations: 9999 (Random sample from a large number)

Number of permuted statistics greater than or equal to R: 0





Table N-49 PERMANOVA test of 'Site' and 'Year' factors for edge habitat of Calna Creek near Hornsby Heights WWTP

Sums of squares type: Type III (partial) Fixed effects sum to zero for mixed terms

Permutation method: Permutation of residuals under a reduced model

Number of permutations: 9999 PERMANOVA table of results

7 27 (10) (100	v / 1 C	011000	1110			Unique
Source	df	SS	MS	Pseudo-F	P(perm)	perms
Site	1	18302	18302	12.125	0.0001	9912
Year	24	52248	2177	1.4423	0.0001	9648
SitexYear	24	41229	1717.9	1.1381	0.0464	9659
Res	46	69434	1509.4			
Total	95	1.8131E+0)5			
Estimates o	f coi	mponents c	of variation			
Source		Estimate	Sq.root			
S(Site)		362.72	19.045			
S(Year)		173.99	13.191			
S(SitexYear)	108.66	10.424			
V(Res)		1509.4	38.851			

Table N-50 PERMANOVA test of 'Site' and 'Year' factors for riffle habitat of Calna Creek near Hornsby Heights WWTP

Sums of squares type: Type III (partial) Fixed effects sum to zero for mixed terms

Permutation method: Permutation of residuals under a reduced model

Number of permutations: 9999 PERMANOVA table of results

						Unique
Source	df	SS	MS	Pseudo-F	P(perm)	perms
Site	1	35302	35302	24.911	0.0001	9930
Year	24	52242	2176.7	1.536	0.0001	9687
SitexYear	24	43221	1800.9	1.2708	0.0023	9680
Res	41	58103	1417.1			
Total	90	1.9243E+0)5			
Estimates of	of coi	mponents c	of variation			
Source		Estimate	Sq.root			
S(Site)		799.68	28.279			
S(Year)		212.92	14.592			
S(SitexYear	r)	215.13	14.667			
V(Res)	-	1417.1	37.645			





Table N-51 PERMDISP test of 'Site' factor for edge habitat of Calna Creek near Hornsby Heights WWTP

DEVIATIONS FROM CENTROID

F: 30.288 df1: 1 df2: 94

P(perm): 0.0001

MEANS AND STANDARD ERRORS
Group Size Average SE
Downstream 48 43.939 0.93465
Upstream 48 37.367 0.74336

Table N-52 PERMDISP test of 'Site' factor for riffle habitat of Calna Creek near Hornsby Heights WWTP

DEVIATIONS FROM CENTROID

F: 2.7061 df1: 1 df2: 89

P(perm): 0.1294

MEANS AND STANDARD ERRORS
Group Size Average SE
Downstream 48 41.739 1.2992
Upstream 43 38.932 1.0705

Table N-53 ANOSIM test of 'Site period' for edge habitat of Calna Creek near Hornsby Heights WWTP



Numbers

Tests for differences between unordered Site period groups

Global Test

Sample statistic (R): 0.299

Significance level of sample statistic: 0.01%

Number of permutations: 9999 (Random sample from a large number)

Number of permuted statistics greater than or equal to R: 0

Pairwise Tests

	K	Significance	Possible	Actual	Number >=
Groups	Statistic	Level %	Permutations	Permutations	Observed
Downstream 1996 to 2018, Downstream 2018 to 2020	-0.078	67.6	194580	9999	6762
Downstream 1996 to 2018, Upstream 1996 to 2018	0.327	0.01	Very large	9999	0
Downstream 1996 to 2018, Upstream 2018 to 2020	0	46.9	194580	9999	4689
Downstream 2018 to 2020, Upstream 1996 to 2018	0.851	0.01	194580	9999	0
Downstream 2018 to 2020, Upstream 2018 to 2020	0.948	2.9	35	35	1
Upstream 1996 to 2018, Upstream 2018 to 2020	0.003	44.8	194580	9999	4478

Cianificance

Table N-54 ANOSIM test of 'Site period' for riffle habitat of Calna Creek near Hornsby Heights WWTP

Tests for differences between unordered Site period groups

Global Test

Sample statistic (R): 0.579

Significance level of sample statistic: 0.01%

Number of permutations: 9999 (Random sample from a large number)

Number of permuted statistics greater than or equal to R: 0

Pairwise Tests

	R	Significance	Possible	Actual	Number >=
Groups	Statistic	Level %	Permutations	Permutations	Observed
Downstream 1996 to 2018, Downstream 2018 to 2020	0.393	1.5	194580	9999	147
Downstream 1996 to 2018, Upstream 1996 to 2018	0.603	0.01	Very large	9999	0
Downstream 1996 to 2018, Upstream 2018 to 2020	0.574	0.07	194580	9999	6
Downstream 2018 to 2020, Upstream 1996 to 2018	0.936	0.02	123410	9999	1
Downstream 2018 to 2020, Upstream 2018 to 2020	0.823	2.9	35	35	1
Upstream 1996 to 2018, Upstream 2018 to 2020	0.162	15.9	123410	9999	1587





Table N-55 Genera subset whose multivariate pattern matches full genera set of the edge habitat Calna Creek near Hornsby Heights WWTP

Subset of 36 (correlation 0.950) genera from edge habitat whose pattern matches that of the full set of 139 genera identified with same subset found on seven runs from 50 random start runs. Each run was based on three randomly selected genera. Genera were:

Chironomidae Chironomus, Physidae Physella, Chironomidae Cricotopus, Chironomidae Cryptochironomus, Chironomidae Dicrotendipes, Coenagrionidae Austroagrion, Dugesiidae Cura, Glossiphoniidae Helobdella, Lumbriculidae Lumbriculus, Lymnaeidae Pseudosuccinea, Naididae Branchiura, Planorbidae Ferrissia, Argiolestidae Austroargiolestes, Chironomidae Microtendipes, Chironomidae Polypedilum, Chironomidae Procladius, Chironomidae Rheocricotopus, Corduliidae Hemicordulia, Tateidae Potamopyrgus, Ceratopogonidae Bezzia, Elmidae Notriolus, Elmidae Simsonia, Gomphidae Austrogomphus, Hydroptilidae Hellyethira, Notonectidae Enithares, Chironomidae Paramerina, Chironomidae Riethia, Corydalidae Archichauliodes, Elmidae Kingolus, Gerridae Tenagogerris, Notonectidae Anisops, Oxidae Oxus, Psephenidae Sclerocyphon, Veliidae Microvelia, Chironomidae Tanytarsus, Stratiomyidae Odontomyia.

Table N-56 Genera subset whose multivariate pattern matches full genera set of the riffle habitat Calna Creek near Hornsby Heights WWTP

Subset of 30 (correlation 0.951) genera from riffle habitat whose pattern matches that of the full set of 105 genera identified with same subset found on 14 runs from 50 random start runs. Each run was based on three randomly selected genera. Genera were:

Chironomidae *Chironomus*, Physidae *Physella*, Chironomidae *Cricotopus*, Dugesiidae *Cura*, Lumbriculidae *Lumbriculus*, Simuliidae *Simulium*, Argiolestidae *Austroargiolestes*, Chironomidae *Polypedilum*, Chironomidae *Procladius*, Chironomidae *Rheocricotopus*, Chironomidae *Rheotanytarsus*, Gelastocoridae *Nerthra*, Hydropsychidae *Cheumatopsyche*, Tateidae *Potamopyrgus*, Ceratopogonidae *Bezzia*, Elmidae *Notriolus*, Elmidae *Simsonia*, Hydroptilidae *Hellyethira*, Aeshnidae *Austroaeschna*, Chironomidae *Parakiefferiella*, Chironomidae *Paratanytarsus*, Corydalidae *Archichauliodes*, Elmidae *Kingolus*, Psephenidae *Sclerocyphon*, Antipodoecidae *Antipodoecia*, Calamoceratidae *Anisocentropus*, Elmidae *Austrolimnius*, Gomphidae *Hemigomphus*, Philopotamidae *Chimarra*, Stratiomyidae *Odontomyia*.








Figure N-39 Shade plot of freshwater macroinvertebrate edge habitat community structure of upstream-downstream sites of Calna Creek near Hornsby Heights WWTP






Figure N-40 Shade plot of freshwater macroinvertebrate riffle habitat community structure of upstream-downstream sites of Calna Creek near Hornsby Heights WWTP





Stream health near West Hornsby WWTP

The SIGNAL-SG plot for Waitara Creek provided an assessment of stream health. This plot was based on macroinvertebrate identification and counting results expressed as SIGNAL-SG scores and allows a visual comparison of data collected from 2019–20 against that collected between 1996 and 2019. This comparison suggests mean stream health of both sites was similar to the preceding 2018-19 year but the range of returned SIGNAL-SG scores was tighter for both sites in 2019-20. This outcome suggested wastewater discharge from West Hornsby WWTP did have a measurable negative impact on stream health during 2019-20 (Figure N-41).

A comparison of the upstream-downstream SIGNAL-SG scores for 2019-20 samples under a t-test returned a significant test outcome (Table N-57) and confirmed the visual trend of little overlap in the range of stream health between upstream and downstream sites in 2019-20.

As a measurable negative impact on downstream stream health was evident, further data analysis was undertaken.




Figure N-41 Stream health of Waitara Creek near West Hornsby WWTP

Table N-57 t-test of upstream-downstream SIGNAL-SG scores of 2019-20 samples from Waitara Creek near West Hornsby WWTP

Method	Variances	DF	t Value	Pr > t	
Pooled	Equal	10	-3.03	0.0127	

Equality of Variances					
Method	Num DF	Den DF	F Value	Pr > F	
Folded F	5	5	1.46	0.6861	



Both edge and pool rock habitats were collected consistently enough at upstreamdownstream sites on the same sampling occasions to allow a multivariate analysis for the monitoring period of 1996 to 2020.

Abutting groups of samples were evident in the three-dimensional nMDS ordination plot of the Waitara Creek edge habitat (Figure N-42). The ordination pattern was confirmed in the corresponding tree diagram (Figure N-44) from classification analysis where the sixth division separated most of the upstream and downstream samples. The pool rock habitat displayed a slightly overlapping pattern, with more recent (2017-2020) downstream samples grouped with other more disparate downstream samples that were generally collected from more later years (Figure N-45).

Shade plot patterns display a smaller set of taxa for each habitat at the downstream site (Figure N-46 and Figure N-47). The BVSTEP routine was used to find a subset of taxa whose multivariate pattern matched that of the full dataset with 34 taxa identified for the edge habitat (Table N-66) and 27 taxa for the pool rock habitat (Table N-67). These subsets of taxa reflect those taxa which formed the main visual patterns in the respective shade plots.

The PERMDISP analysis indicated a significantly different pattern of dispersion (spacing between same samples) for the upstream and downstream sites of the edge and pool rock habitats (Table N-62 and Table N-63). This suggests the variability in taxonomic composition of samples collected over time was different for upstream and downstream sites through the period 1996 to 2020. This outcome suggests subsequent edge and pool rock habitat results of ANOSIM tests are describing both the variability in taxonomic composition of samples over time as well as community composition variability between the upstream and downstream sites.

An ANOSIM test was run on the factor 'Site'. The returned R-values were at a mid-range level for edge (Table N-58) and at a low-range level for pool rock (Table N-59). These R-value results suggest site specific assemblages were more distinguishable for the edge habitat and less distinguishable for the pool rock habitat.

To further explore community structure, hypothesis testing was conducted with a PERMANOVA model. This model comprised the fixed factors 'Site' and 'Year' with 'Year' representing samples collected between 1996 and 2020 and 'Site' having two levels, upstream and downstream. A statistically non-significant 'Site x Year' interaction was returned for both the edge and pool rock habitats (Table N-60 and Table N-61). These non-significant results allowed us to view the 'Site' and 'Year' results. Significant results were returned for 'Site' and 'Year' for both habitats.

A second run of ANOSIM based on 'Site-Period' sample groups displayed in the above ordination plots (Figure N-42 and Figure N-43) returned a significant global mid-range R-value (0.46) for the edge habitat. In the resulting pairwise comparisons, three of the six tests returned significant R-values at a mid to high level (Table N-64). A slightly lower mid-range global R-value of 0.386 was returned for the pool rock habitat. A corresponding pairwise test for both the edge and pool rock habitats returned high level significant R-value for the comparison of 2017 to 2020 downstream site to the 1996 to 2017 upstream site time periods (Table N-65). This test outcome likely reflects disturbance by wastewater discharge as it is above the 0.66 R-value determined by Besley and Chessman (2008) that represents natural habitat differences between sites on the same stream.

SIGNAL-SG and multivariate testing outcomes suggest downstream community structure in Waitara Creek was altered by wastewater discharge from West Hornsby WWTP in the more recent period.








Figure N-42 Dimensions 1 and 2 of three-dimensional ordination plot of freshwater macroinvertebrate edge habitat community structure of upstream-downstream sites of Waitara Creek near West Hornsby WWTP




Figure N-43 Dimensions 1 and 2 of three-dimensional ordination plot of freshwater macroinvertebrate pool rock habitat community structure of upstream-downstream sites of Waitara Creek near West Hornsby WWTP








Figure N-44 Tree diagram of freshwater macroinvertebrate edge habitat community structure of upstream-downstream sites of Waitara Creek near West Hornsby WWTP




Figure N-45 Tree diagram of freshwater macroinvertebrate pool rock habitat community structure of upstream-downstream sites of Waitara Creek near West Hornsby WWTP





Table N-58 ANOSIM test of 'Site' factor for edge habitat of Waitara Creek near West Hornsby WWTP

Analysis of Similarities Site levels Downstream Upstream

Tests for differences between unordered Site groups Global Test Sample statistic (R): 0.502

Significance level of sample statistic: 0.01%

Number of permutations: 9999 (Random sample from a large number)

Table N-59 ANOSIM test of 'Site' factor for pool rock habitat of Waitara Creek near West Hornsby WWTP

Analysis of Similarities Site levels Downstream Upstream

Tests for differences between unordered Site groups Global Test Sample statistic (R): 0.319

Significance level of sample statistic: 0.01%

Number of permutations: 9999 (Random sample from a large number)





Table N-60 PERMANOVA test of 'Site' and 'Year' factors for edge habitat of Waitara Creek near West Hornsby WWTP

Sums of squares type: Type III (partial) Fixed effects sum to zero for mixed terms

Permutation method: Permutation of residuals under a reduced model

Number of permutations: 9999

PERMANOVA table of results

						Unique
Source	df	SS	MS	Pseudo-F	P(perm)	perms
Site	1	24278	24278	19.54	0.0001	9923
Year	24	55426	2309.4	1.8588	0.0001	9657
SitexYear	24	33547	1397.8	1.125	0.0717	9676
Res	48	59637	1242.4			
Total	97	1.7377E+05				

Estimates of components of variation

Source	Estimate	Sq.root
S(Site)	491.42	22.168
S(Year)	272.54	16.509
S(SitexYear)	79.358	8.9083
V(Res)	1242.4	35.248

Table N-61 PERMANOVA test of 'Site' and 'Year' factors for pool rock habitat of Waitara Creek near West Hornsby WWTP

Sums of squares type: Type III (partial) Fixed effects sum to zero for mixed terms

Permutation method: Permutation of residuals under a reduced model

Number of permutations: 9999

PERMANOVA table of results

- El tim il to tri table of roodite							
						Unique	
Source	df	SS	MS	Pseudo-F	P(perm)	perms	
Site	1	15456	15456	12.956	0.0001	9918	
Year	24	51862	2160.9	1.8114	0.0001	9699	
SitexYear	23	30808	1339.5	1.1228	0.1131	9716	
Res	42	50106	1193				
Total	90	1.5306E+05					

Estimates of components of variation

	oompomome or	ranacioni
Source	Estimate	Sq.root
S(Site)	342.54	18.508
S(Year)	272.75	16.515
S(SitexYear)	80.082	8.9488
V(Res)	1193	34.54





Table N-62 PERMDISP test of 'Site' factor for edge habitat of Waitara Creek near West Hornsby WWTP

DEVIATIONS FROM CENTROID

F: 32.332 df1: 1 df2: 96

P(perm): 0.0001

MEANS AND STANDARD ERRORS

 Group
 Size
 Average
 SE

 Downstream
 49
 41.752
 0.9345

 Upstream
 49
 34.939
 0.74991

Table N-63 PERMDISP test of 'Site' factor for pool rock habitat of Waitara Creek near West Hornsby WWTP

DEVIATIONS FROM CENTROID

F: 18.989 df1: 1 df2: 89

P(perm): 0.0002

MEANS AND STANDARD ERRORS

Group Size Average SE Downstream 48 40.739 0.96506 Upstream 43 34.716 0.98587





Table N-64 ANOSIM test of 'Site period' factor for edge habitat of Waitara Creek near West Hornsby WWTP

Global Test

Sample statistic (R): 0.46

Significance level of sample statistic: 0.01%

Number of permutations: 9999 (Random sample from a large number)

Number of permuted statistics greater than or equal to R: 0

Pairwise Tests

R	Significance	Possible	Actual	Number >=
Statistic	Level %	Permutations	Permutations	Observed
0.241	3.8	13983816	9999	381
0.512	0.01	Very large	9999	0
0.233	3.5	13983816	9999	353
0.913	0.01	13983816	9999	0
0.489	0.2	462	462	1
0.04	36	13983816	9999	3603
	0.241 0.512 0.233 0.913 0.489	Statistic Level % 0.241 3.8 0.512 0.01 0.233 3.5 0.913 0.01 0.489 0.2	Statistic Level % Permutations 0.241 3.8 13983816 0.512 0.01 Very large 0.233 3.5 13983816 0.913 0.01 13983816 0.489 0.2 462	Statistic Level % Permutations Permutations 0.241 3.8 13983816 9999 0.512 0.01 Very large 9999 0.233 3.5 13983816 9999 0.913 0.01 13983816 9999 0.489 0.2 462 462

Table N-65 ANOSIM test of 'Site period' factor for pool rock habitat of Waitara Creek near West Hornsby WWTP

Global Test

Sample statistic (R): 0.386

Significance level of sample statistic: 0.01%

Number of permutations: 9999 (Random sample from a large number)

Number of permuted statistics greater than or equal to R: 0

Pairwise Tests

	R	Significance	Possible	Actual	Number >=
Groups	Statistic	Level %	Permutations	Permutations	Observed
Downstream 1996 to 2017, Downstream 2017 to 2020	0.315	0.6	12271512	9999	54
Downstream 1996 to 2017, Upstream 1996 to 2017	0.345	0.01	Very large	9999	0
Downstream 1996 to 2017, Upstream 2017 to 2020	0.186	10.7	163185	9999	1066
Downstream 2017 to 2020, Upstream 1996 to 2017	0.867	0.01	8145060	9999	0
Downstream 2017 to 2020, Upstream 2017 to 2020	0.524	0.5	210	210	1
Upstream 1996 to 2017, Upstream 2017 to 2020	0.335	1.8	123410	9999	182



00

Table N-66 Genera subset whose multivariate pattern matches full genera set of the edge habitat of Waitara Creek near West Hornsby WWTP

Subset of 34 (correlation 0.952) genera from edge habitat whose pattern matches that of the full set of 142 genera identified with same subset found on 27 runs from 50 random start runs. Each run was based on three randomly selected genera. Genera were:

Tateidae *Posticobia*, Chironomidae *Chironomus*, Erpobdellidae *Vivabdella*, Physidae *Physella*, Planorbidae *Helicorbis*, Chironomidae *Cricotopus*, Chironomidae *Dicrotendipes*, Dugesiidae *Cura*, Glossiphoniidae *Helobdella*, Lumbriculidae *Lumbriculus*, Lymnaeidae *Austropeplea*, Argiolestidae *Austroargiolestes*, Chironomidae *Polypedilum*, Chironomidae *Procladius*, Chironomidae *Rheocricotopus*, Corduliidae *Hemicordulia*, Hydrophilidae *Enochrus*, Hydropsychidae *Cheumatopsyche*, Isostictidae *Rhadinosticta*, Libellulidae *Nannophlebia*, Sphaeriidae *Musculium*, Tateidae *Potamopyrgus*, Ceratopogonidae *Bezzia*, Elmidae *Simsonia*, Gomphidae *Austrogomphus*, Hydroptilidae *Hellyethira*, Notonectidae *Enithares*, Chironomidae *Paramerina*, Elmidae *Kingolus*, Gerridae *Tenagogerris*, Oxidae *Oxus*, Psephenidae *Sclerocyphon*, Veliidae *Microvelia*, Stratiomyidae *Odontomyia*.

Table N-67 Genera subset whose multivariate pattern matches full genera set of the pool rock habitat of Waitara Creek near West Hornsby WWTP

Subset of 27 (correlation 0.952) genera from pool rock habitat whose pattern matches that of the full set of 79 genera identified with same subset found on 10 runs from 50 random start runs. Each run was based on three randomly selected genera. Genera were:

Tateidae *Posticobia*, Erpobdellidae *Vivabdella*, Physidae *Physella*, Planorbidae *Helicorbis*, Chironomidae *Cricotopus*, Chironomidae *Dicrotendipes*, Dugesiidae *Cura*, Glossiphoniidae *Helobdella*, Lumbriculidae *Lumbriculus*, Naididae *Branchiura*, Naididae *Nais*, Argiolestidae *Austroargiolestes*, Chironomidae *Cladotanytarsus*, Chironomidae *Polypedilum*, Chironomidae *Procladius*, Chironomidae *Rheocricotopus*, Glossiphoniidae *Alboglossiphonia*, Hydropsychidae *Cheumatopsyche*, Isostictidae *Rhadinosticta*, Tateidae *Potamopyrgus*, Ecnomidae *Ecnomus*, Elmidae *Notriolus*, Elmidae *Simsonia*, Hydroptilidae *Hellyethira*, Corydalidae *Archichauliodes*, Elmidae *Kingolus*, Psephenidae *Sclerocyphon*.








Figure N-46 Shade plot of freshwater macroinvertebrate edge habitat community structure of upstream-downstream sites of Waitara Creek near West Hornsby WWTP





Figure N-47 Shade plot of freshwater macroinvertebrate pool rock habitat community structure of upstream-downstream sites of Waitara Creek near West Hornsby WWTP





Other sites of the Hawkesbury-Nepean River system that are only assessed under the macroinvertebrate indicator and not directly related to WWTP discharges follow.


Cattai Creek at Annangrove Road, d/s of both Rouse Hill and Castle Hill plants



The range of stream health recorded over each period is represented by length of line

No samples were taken at NC5 in 2019-20 due to safety concerns associated with illegal dumping of asbestos near the sampling site.

Hawkesbury River at Windsor Bridge, u/s South Creek




The range of stream health recorded over each period is represented by length of line

Mean stream health in 2019-20 was within the range recorded over the period back to 1995.







Hawkesbury River at Wilberforce



The range of stream health recorded over each period is represented by length of line

Mean stream health in 2019-20 was within the range recorded over the period back to 1995.





The range of stream health recorded over each period is represented by length of line

Mean stream health in 2019-20 was within the range recorded over longer period back to 1995.





Appendix 0 : Stream health of other Sydney rivers

Sites monitored for the macroinvertebrate indicator in freshwater streams assessed the general condition of stream health in urban areas. Among these, four are control sites located upstream of any likely impact from urban areas. Three out of the four urban sites are situated in areas just upstream of estuarine limits of the Parramatta River (PJPR), Lane Cove River (PJLC) and Georges River (GR22). The fourth urban site is situated about 5 km further up in the Georges River (GR23). The control sites were Lynch's Creek (N451) a tributary of Hawkesbury-Nepean River, Hacking River at McKell Avenue in Royal National Park (PH22), the upper Georges River system at O'Hares Creek (GE510) and Georges River at Ingleburn Reserve (GR24).

Results from 2019-20 indicate stream health for one of the four control sites (N451) was typical of natural water quality in bushland areas that do not receive urban stormwater runoff or sewer overflows (Figure O-1 and Figure O-2). The mean stream health for the Hacking River at Mckell Avenue fell within the 'mild water pollution' category (Figure O-1). Additionally, mean stream health for the Georges River at Ingleburn Reserve (GR24 - control site) and O'Hares Creek were within the 'mild water pollution' category (Figure O-2).

Urban sites within the Port Jackson Rivers upstream of Lane Cove Weir (PJLC) and Parramatta Weir (PJPR) fell around the boundary of the 'mild and moderate water pollution' category (Figure O-1). Additionally, urban sites within the Georges River for 2019-20 fell within the 'mild water pollution' category (Figure O-2). These results were near the upper part of the range of stream health that has been recorded for these urban sites over the previous 1995 to 2019 period.

Results from these test sites represent the ambient condition of the combined impact of urban stormwater runoff and sewer overflows. These two influences cannot be teased apart. Pilot studies are being designed under Sydney Water's wet weather overflow program to model a stormwater control condition, which may then better define the impact of sewer overflows on urban streams than these current single site assessments.








Figure O-1 Stream health of Lane Cove and Parramatta rivers in comparison to control sites





Figure O-2 Stream health of lower freshwater Georges River sites compared to control sites in the upper Georges River system







Publication Number: SW135 12/20 Sydney Water. All rights reserved.

